arXiv:1605.00476v1 [math.CA] 2 May 2016

BOUNDEDNESS OF BI-PARAMETER LITTLEWOOD-PALEY OPERATORS ON PRODUCT HARDY SPACE

ZHENGYANG LI AND QINGYING XUE

Abstract. Let $n_1, n_2 \geq 1, \lambda_1 > 1$ and $\lambda_2 > 1$. For any $x = (x_1, x_2) \in \mathbb{R}^n \times \mathbb{R}^m$, let g and $g^*_{\vec{\lambda}}$ be the bi-parameter Littlewood-Paley square functions defined by

$$g(f)(x) = \Big(\int_0^\infty \int_0^\infty |\theta_{t_1,t_2} f(x_1, x_2)|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \Big)^{1/2}, \quad \text{and}$$

$$g^*_{\vec{\lambda}}(f)(x) = \Big(\iint_{\mathbb{R}^{m+1}_+} \iint_{\mathbb{R}^{n+1}_+} \prod_{i=1}^2 \Big(\frac{t_1}{t_i + |x_i - y_i|} \Big)^{n_i \lambda_i} |\theta_{t_1,t_2} f(y_1, y_2)|^2 \frac{dy_1 dt_1}{t_1^{n+1}} \frac{dy_2 dt_2}{t_2^{m+1}} \Big)^{1/2},$$

where $\theta_{t_1,t_2} f(x_1, x_2) = \iint_{\mathbb{R}^n \times \mathbb{R}^m} s_{t_1,t_2}(x_1, x_2, y_1, y_2) f(y_1, y_2) dy_1 dy_2$. It is known that the L^2 boundedness of bi-parameter g and $g^*_{\vec{\lambda}}$ have been established recently by Martikainen, and Cao, Xue, respectively. In this paper, under certain structure conditions assumed on the kernel s_{t_1,t_2}, we show that both g and $g^*_{\vec{\lambda}}$ are bounded from product Hardy space $H^1(\mathbb{R}^n \times \mathbb{R}^m)$ to $L^1(\mathbb{R}^n \times \mathbb{R}^m)$. As consequences, the L^p boundedness of g and $g^*_{\vec{\lambda}}$ will be obtained for $1 < p < 2$.

1. Introduction

1.1. **Bi-parameter Littlewood-Paley operators.** The study of multi-parameter operators originated in the famous works of Fefferman and Stein [6] on bi-parameter singular integral operators. Later on, Journé [9] peresented a multi-parameter version of $T1$ theorem on product spaces. Subsequently, Pott and Villarroya [12] formulated a new type of $T1$ theorem. Recently, a bi-parameter representation of singular integrals in expression of the dyadic shifts was given by Martikainen [10], which extended the famous result of Hytönen [7] for one-parameter case. Moreover, a bi-parameter version of $T1$ theorem in spaces of non-homogeneous type was demonstrated by Hytönen and Martikainen [8].

Still more recently, Martikainen [18] studied a class of bi-parameter square function defined as follows:

Definition 1.1. ([18]) For any $x = (x_1, x_2) \in \mathbb{R}^n \times \mathbb{R}^m$, the bi-parameter Littlewood-Paley square function g is defined by

$$g(f)(x) := \Big(\int_0^\infty \int_0^\infty |\theta_{t_1,t_2} f(x_1, x_2)|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \Big)^{1/2},$$

Date: May 2, 2016.

Key words and phrases. Bi-parameter square function; Bi-parameter g^*_λ-function; Product H^1 space; Rectangle atomic decomposition.

The second author was supported partly by NSFC (No. 11471041), the Fundamental Research Funds for the Central Universities (NO. 2014KJJCA10) and NCET-13-0065.

Corresponding author: Qingying Xue Email: qyxue@bnu.edu.cn.

where $\theta_{t_1,t_2}f(x_1,x_2)=\iint_{\mathbb{R}^n\times\mathbb{R}^m} s_{t_1,t_2}(x_1,x_2,y_1,y_2)f(y_1,y_2)dy_1\,dy_2$.

In [18], the kernels s_{t_1,t_2} were assumed to satisfy a natural size estimate, a Hölder estimates and two symmetric mixed Hölder and size estimates, they were also assumed to satisfy certain mixed Carleson and size estimates, mixed Carleson and Hölder estimates and a bi-parameter Carleson condition. Under these assumptions, the author of [18] established the L^2 boundedness of g.

Later on, by modifying and adding new ingredients in the assumptions, Cao and Xue [1] introduced and studied the L^2 boundedness of the following bi-parameter Littlewood-Paley g^*_λ-function:

Definition 1.2. ([1]) Let $\lambda_1,\lambda_2>1$, for any $x=(x_1,x_2)\in\mathbb{R}^n\times\mathbb{R}^m$, the bi-parameter Littlewood-Paley g^*_λ-function is defined by

$$g^*_\lambda(f)(x):=\Big(\iint_{\mathbb{R}^{m+1}_+}\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\iint_{\mathbb{R}^{n+1}_+}\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}\times|\vartheta_{t_1,t_2}f(y_1,y_2)|^2\frac{dy_1dt_1}{t_1^{n+1}}\frac{dy_2dt_2}{t_2^{m+1}}\Big)^{1/2},$$

where $\vartheta_{t_1,t_2}f(y_1,y_2)=\iint_{\mathbb{R}^n\times\mathbb{R}^m}K_{t_1,t_2}(y_1,y_2,z_1,z_2)f(z_1,z_2)dz_1\,dz_2$.

Till now, the only known result about g-function and g^*_λ-functionis is the L^2 boundedness. Therefore, it leaves several questions open, such as, it is quite natural to ask if the L^p boundedness and H^1 to L^1 boundedness are true or not. In this paper, our objects of investigation are the bi-parameter Littlewood-Paley g-function and g^*_λ-function. We are mainly concerned with the boundedness of these two operators on product Hardy spaces (as consequences, the L^p boundedness will be obtained for $1<p<2$). The introduction of product Hardy spaces will be given in the next subsection.

1.2. **Product Hardy space.** The product Hardy space was first introduced and studied by Malliavin and Malliavin [17] in 1977, and systematically studied by Gundy and Stein [13] in 1979. Later on, Chang and R. Fefferman [3] established the atomic decomposition of $H^p(\mathbb{R}^2_+\times\mathbb{R}^2_+)$, which is more complicated than the classical $H^p(\mathbb{R}^n)$. By using the rectangle atomic decomposition of $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ (Theorem A below) and a geometric covering lemma due to Journé[16], R. Fefferman [5] established the $(H^p(\mathbb{R}^n\times\mathbb{R}^m),L^p(\mathbb{R}^n\times\mathbb{R}^m))$ boundedness of Journé's product singular integrals. For more work about the boundedness of operators on product Hardy space, one may refer to [4], [14], [15], [20].

In order to state some known results, we need to introduce two more definitions.

Definition 1.3 ($H^p(\mathbb{R}^n\times\mathbb{R}^m)$ **atom**, [2])**.** Let $a(x_1,x_2)$ be a function supported in an open set $\Omega\subset\mathbb{R}^n\times\mathbb{R}^m$ with finite measure. $a(x_1,x_2)$ is said to be an $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ atom if it satisfies the following condition:

(i) $\|a\|_2\le|\Omega|^{1/2-1/p}$.

(ii) a can further be decomposed as $a(x_1,x_2)=\sum_{R\in\mathcal{M}(\Omega)}a_R(x_1,x_2)$, where a_R are supported on the double of $R=I\times J$ (I a dyadic cube in $\mathbb{R}^n$, J a dyadic cube

in $\mathbb{R}^m$) and $\mathcal{M}(\Omega)$ is the collection of all maximal dyadic rectangles contained in Ω. Moreover, $\big\{\sum_{R\in\mathcal{M}(\Omega)}\|a_R\|_2^2\big\}^{1/2}\le|\Omega|^{1/2-1/p}$.

(iii) $\int_{2I}a_R(x_1,x_2)x_1^\alpha dx_1=0$ for all $x_2\in\mathbb{R}^m$, $0\le|\alpha|\le N_{p,n}$,
$\int_{2J}a_R(x_1,x_2)x_2^\alpha dx_2=0$ for all $x_1\in\mathbb{R}^n$, $0\le|\beta|\le N_{p,m}$,
where $N_{p,n}$ and $N_{p,m}$ is a large integer depending on p and n,

Definition 1.4 ($H^p(\mathbb{R}^n\times\mathbb{R}^m)$ **rectangle atom**, [2])**.** Let function $a(x_1,x_2)$ be supported on a rectangle $R=I\times J$, where I is a cube in $\mathbb{R}^n$ and J is a cube in $\mathbb{R}^m$, respectively, $a(x_1,x_2)$ is called an $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom provided

(i) $\|a\|_2\le|R|^{1/2-1/p}$,
(ii) $\int_I a(x_1,x_2)x_1^\alpha dx_1=0$ for all $x_2\in\mathbb{R}^m$, $0\le|\alpha|\le N_{n,p}$,
(iii) $\int_J a(x_1,x_2)x_2^\beta dx_2=0$ for all $x_1\in\mathbb{R}^n$, $0\le|\beta|\le N_{m,p}$.

Chang and R. Fefferman [2] gave the following atomic decomposition of $H^p(\mathbb{R}^n\times\mathbb{R}^m)$.

Theorem A. ([2]) *A distribution $f\in H^p(\mathbb{R}^n\times\mathbb{R}^m)$ if and only if $f=\sum_j\lambda_j a_j$, where a_j are $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ atoms, $\sum_j|\lambda_j|^p<\infty$, and the series converges in the distribution sense. Moreover, $\|f\|_{H^p}^p$ is equivalent to $\inf\{\sum_j|\lambda_j|^p$: for all $f=\sum_j\lambda_j a_j\}$.*

Since the support of $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ atom is an open set, the decomposition of $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ cannot be carried out to demonstrate the H^p boundedness of linear or sublinear operators directly as the classical cases. However, it was quite surprising that R. Fefferman [5] obtained the following nice result,

Theorem B. ([5]) *Suppose that T is a bounded linear operator on $L^2(\mathbb{R}^n\times\mathbb{R}^m)$. Suppose further that if a is an $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom $(0<p\le1)$ supported on R, we have*

$$\iint_{\mathbb{R}^n\times\mathbb{R}^m\setminus\widetilde{R}_\gamma}|T(a)(x_1,x_2)|^p dx_1dx_2\le C\gamma^{-\delta},\qquad \text{for all } \gamma\ge2 \text{ and some fixed } \delta>0,$$

where $\widetilde{R}_\gamma$ denotes the γ fold enlargement of R. Then T is a bounded operator from $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ to $L^p(\mathbb{R}^n\times\mathbb{R}^m)$.

1.3. **Assumptions and Main result.**

In order to state our main results, we first present the following assumptions:
Assumption 1 (Standard estimates of s_{t_1,t_2}). The kernel $s_{t_1,t_2}:(\mathbb{R}^n\times\mathbb{R}^m)\times(\mathbb{R}^n\times\mathbb{R}^m)\to\mathbb{C}$ is assumed to satisfy the following estimates:

(1) Size condition :

$$|s_{t_1,t_2}(x,y)|\lesssim\frac{t_1^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{t_2^\beta}{(t_2+|x_2-y_2|)^{m+\beta}}.$$

(2) Hölder condition :

$$|s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,y_2'))-s_{t_1,t_2}(x,(y_1',y_2))+s_{t_1,t_2}(x,y')|\lesssim\frac{|y_1-y_1'|^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{|y_2-y_2'|^\beta}{(t_2+|x_2-y_2|)^{m+\beta}},$$

whenever $|y_1-y_1'|<t_1/2$ and $|y_2-y_2'|<t_2/2$.

(3) Mixed Hölder and size conditions :

$$|s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,y_2'))|\lesssim \frac{t_1^{\alpha}}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{|y_2-y_2'|^{\beta}}{(t_2+|x_2-y_2|)^{m+\beta}},$$

whenever $|y_2-y_2'|<t_2/2$ and

$$|s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1',y_2))|\lesssim \frac{|y_1-y_1'|^{\alpha}}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{t_2^{\beta}}{(t_2+|x_2-y_2|)^{m+\beta}},$$

whenever $|y_1-y_1'|<t_1/2$.

Assumption 2 (Carleson condition $\times$ Standard estimates of s_{t_1,t_2}). If $I\subset\mathbb{R}^n$ is a cube with side length $\ell(I)$, we define the associated Carleson box by $\widehat{I}=I\times(0,\ell(I))$. We assume the following conditions : For every cube $I\subset\mathbb{R}^n$ and $J\subset\mathbb{R}^m$, there holds that

(1) Combinations of Carleson and size conditions :

$$\Big(\iint_{\widehat{I}}\Big|\int_I s_{t_1,t_2}(x_1,x_2,y_1,y_2)dy_1\Big|^2\frac{dx_1dt_1}{t_1}\Big)^{1/2}\lesssim |I|^{1/2}\frac{t_2^{\beta}}{(t_2+|x_2-y_2|)^{m+\beta}}$$

and

$$\Big(\iint_{\widehat{J}}\Big|\int_J s_{t_1,t_2}(x_1,x_2,y_1,y_2)dy_2\Big|^2\frac{dx_2dt_2}{t_2}\Big)^{1/2}\lesssim \frac{t_1^{\alpha}}{(t_1+|x_1-y_1|)^{m+\alpha}}|J|^{1/2}.$$

(2) Combinations of Carleson and Hölder conditions :

$$\Big(\iint_{\widehat{I}}\Big|\int_I[s_{t_1,t_2}(x_1,x_2,y_1,y_2)-s_{t_1,t_2}(x_1,x_2,y_1,y_2')]dy_1\Big|^2\frac{dx_1dt_1}{t_1}\Big)^{1/2}\lesssim |I|^{1/2}\frac{|y_2-y_2'|^{\beta}}{(t_2+|x_2-y_2|)^{m+\beta}},$$

whenever $|y_2-y_2'|<t_2/2$. And

$$\Big(\iint_{\widehat{J}}\Big|\int_J[s_{t_1,t_2}(x_1,x_2,y_1,y_2)-s_{t_1,t_2}(x_1,x_2,y_1',y_2)]dy_2\Big|^2\frac{dx_2dt_2}{t_2}\Big)^{1/2}\lesssim |J|^{1/2}\frac{|y_1-y_1'|^{\alpha}}{(t_1+|x_1-y_1|)^{m+\alpha}},$$

whenever $|y_1-y_1'|<t_1/2$.

Assumption 2′ (Strong Carleson condition $\times$ Standard estimates of s_{t_1,t_2}). If $I\subset\mathbb{R}^n$ is a cube with side length $\ell(I)$, we define the associated Carleson box by $\widehat{I}=I\times(0,\ell(I))$. We assume the following conditions : For every cube $I\subset\mathbb{R}^n$ and $J\subset\mathbb{R}^m$, there holds that

(1) Combinations of Carleson and size conditions :

$$\Big(\iint_{\widehat{I}}\int_I|s_{t_1,t_2}(x_1,x_2,y_1,y_2)|^2dy_1\frac{dx_1dt_1}{t_1}\Big)^{1/2}\lesssim\frac{t_2^{\beta}}{(t_2+|x_2-y_2|)^{m+\beta}}$$

and

$$\Big(\iint_{\widehat{J}}\int_J|s_{t_1,t_2}(x_1,x_2,y_1,y_2)|^2dy_2\frac{dx_2dt_2}{t_2}\Big)^{1/2}\lesssim\frac{t_1^{\beta}}{(t_1+|x_1-y_1|)^{m+\alpha}}.$$

(2) Combinations of Carleson and Hölder conditions :

$$\Big(\iint_{\widehat{I}}\int_I |s_{t_1,t_2}(x_1,x_2,y_1,y_2)-s_{t_1,t_2}(x_1,x_2,y_1,y_2')|^2 dy_1\frac{dx_1dt_1}{t_1}\Big)^{1/2} \lesssim \frac{|y_2-y_2'|^\beta}{(t_2+|x_2-y_2|)^{m+\beta}},$$

whenever $|y_2-y_2'|<t_2/2$. And

$$\Big(\iint_{\widehat{J}}\int_J |s_{t_1,t_2}(x_1,x_2,y_1,y_2)-s_{t_1,t_2}(x_1,x_2,y_1',y_2)|^2 dy_2\frac{dx_2dt_2}{t_2}\Big)^{1/2} \lesssim \frac{|y_1-y_1'|^\alpha}{(t_1+|x_1-y_1|)^{m+\alpha}},$$

whenever $|y_1-y_1'|<t_1/2$.

Assumption 3 (Bi-parameter Carleson condition of s_{t_1,t_2}). Let $\mathcal{D}=\mathcal{D}_n\times\mathcal{D}_m$, where $\mathcal{D}_n$ is a dyadic grid in $\mathbb{R}^n$ and $\mathcal{D}_m$ is a dyadic grid in $\mathbb{R}^m$. For $I\in\mathcal{D}_n$, let $W_I=I\times(\ell(I)/2,\ell(I))$ be the associated Whitney region. Denote

$$C_{IJ}^{\mathcal{D}}=\iint_{W_J}\iint_{W_I}|\theta_{t_1,t_2}\mathbf{1}(y_1,y_2)|^2\frac{dx_1dt_1}{t_1}\frac{dx_2dt_2}{t_2}.$$

We assume the following *bi-parameter Carleson condition*: For every $\mathcal{D}=\mathcal{D}_n\times\mathcal{D}_m$ there holds that

$$\sum_{\substack{I\times J\in\mathcal{D}\\ I\times J\subset\Omega}} C_{IJ}^{\mathcal{D}}\lesssim|\Omega| \tag{1.1}$$

for all sets $\Omega\subset\mathbb{R}^n\times\mathbb{R}^m$ such that $|\Omega|<\infty$ and such that for every $x\in\Omega$ there exists $I\times J\in\mathcal{D}$ so that $x\in I\times J\subset\Omega$.

Assumption 4 (Standard estimates of K_{t_1,t_2}). The kernel $K_{t_1,t_2}:(\mathbb{R}^n\times\mathbb{R}^m)\times(\mathbb{R}^n\times\mathbb{R}^m)\to\mathbb{C}$ is assumed to satisfy the following estimates:

(1) Size condition :

$$|K_{t_1,t_2}(x,y)|\lesssim\frac{t_1^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{t_2^\beta}{(t_2+|x_2-y_2|)^{m+\beta}}.$$

(2) Hölder condition :

$$|K_{t_1,t_2}(x,y)-K_{t_1,t_2}(x,(y_1,y_2'))-K_{t_1,t_2}(x,(y_1',y_2))+K_{t_1,t_2}(x,y')| \lesssim\frac{|y_1-y_1'|^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{|y_2-y_2'|^\beta}{(t_2+|x_2-y_2|)^{m+\beta}},$$

whenever $|y_1-y_1'|<t_1/2$ and $|y_2-y_2'|<t_2/2$.

(3) Mixed Hölder and size conditions :

$$|K_{t_1,t_2}(x,y)-K_{t_1,t_2}(x,(y_1,y_2'))|\lesssim\frac{t_1^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{|y_2-y_2'|^\beta}{(t_2+|x_2-y_2|)^{m+\beta}},$$

whenever $|y_2 - y_2'| < t_2/2$ and

$$|K_{t_1,t_2}(x,y) - K_{t_1,t_2}(x,(y_1',y_2))| \lesssim \frac{|y_1-y_1'|^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}} \frac{t_2^\beta}{(t_2+|x_2-y_2|)^{m+\beta}},$$

whenever $|y_1 - y_1'| < t_1/2$.

Assumption 5 (Carleson condition $\times$ Standard estimates of K_{t_1,t_2}). If $I \subset \mathbb{R}^n$ is a cube with side length $\ell(I)$, we define the associated Carleson box by $\widehat{I} = I \times (0, \ell(I))$. We assume the following conditions : For every cube $I \subset \mathbb{R}^n$ and $J \subset \mathbb{R}^m$, there holds that

(1) Combinations of Carleson and size conditions :

$$\Big(\iint_{\widehat{I}} \int_{\mathbb{R}^n} \Big| \int_I K_{t_1,t_2}(x-y,z_1,z_2)dz_1 \Big|^2 \Big(\frac{t_1}{t_1+|y_1|} \Big)^{n\lambda_1} \frac{dy_1}{t_1^n} \frac{dx_1 dt_1}{t_1} \Big)^{1/2} \lesssim |I|^{1/2} \frac{t_2^\beta}{(t_2+|x_2-y_2-z_2|)^{m+\beta}}$$

and

$$\Big(\iint_{\widehat{J}} \int_{\mathbb{R}^m} \Big| \int_J K_{t_1,t_2}(x-y,z_1,z_2)dz_2 \Big|^2 \Big(\frac{t_2}{t_2+|y_2|} \Big)^{n\lambda_1} \frac{dy_2}{t_2^n} \frac{dx_2 dt_2}{t_2} \Big)^{1/2} \lesssim |J|^{1/2} \frac{t_1^\alpha}{(t_1+|x_1-y_1-z_1|)^{n+\alpha}}.$$

(2) Combinations of Carleson and Hölder conditions :

$$\Big(\iint_{\widehat{I}} \int_{\mathbb{R}^n} \Big| \int_I [K_{t_1,t_2}(x-y,z_1,z_2) - K_{t_1,t_2}(x-y,z_1,z_2')]dz_1 \Big|^2 \Big(\frac{t_1}{t_1+|y_1|} \Big)^{n\lambda_1} \frac{dy_1}{t_1^n} \frac{dx_1 dt_1}{t_1} \Big)^{\frac{1}{2}} \lesssim |I|^{1/2} \frac{|z_2-z_2'|^\beta}{(t_2+|x_2-y_2-z_2|)^{m+\beta}},$$

whenever $|z_2 - z_2'| < t_2/2$. And

$$\Big(\iint_{\widehat{J}} \int_{\mathbb{R}^m} \Big| \int_J [K_{t_1,t_2}(x-y,z_1,z_2) - K_{t_1,t_2}(x-y,z_1',z_2)]dz_2 \Big|^2 \Big(\frac{t_2}{t_2+|y_2|} \Big)^{m\lambda_2} \frac{dy_2}{t_2^n} \frac{dx_2 dt_2}{t_2} \Big)^{\frac{1}{2}} \lesssim |J|^{1/2} \frac{|z_1-z_1'|^\alpha}{(t_1+|x_1-y_1-z_1|)^{n+\alpha}},$$

whenever $|z_1 - z_1'| < t_1/2$.

Assumption 5′ (Strong Carleson condition $\times$ Standard estimates of K_{t_1,t_2}). If $I \subset \mathbb{R}^n$ is a cube with side length $\ell(I)$, we define the associated Carleson box by $\widehat{I} = I \times (0, \ell(I))$. We assume the following conditions : For every cube $I \subset \mathbb{R}^n$ and $J \subset \mathbb{R}^m$, there holds that

(1) Combinations of Carleson and size conditions :

$$\Big(\iint_{\widehat{I}} \int_{\mathbb{R}^n} \int_I |K_{t_1,t_2}(x-y,z_1,z_2)|^2 \Big(\frac{t_1}{t_1+|y_1|} \Big)^{n\lambda_1} \frac{dz_1 dy_1}{t_1^n} \frac{dx_1 dt_1}{t_1} \Big)^{1/2} \lesssim \frac{t_2^\beta}{(t_2+|x_2-y_2-z_2|)^{m+\beta}}$$

and

$$\Big(\iint_{\widehat{J}}\int_{\mathbb{R}^m}\int_J |K_{t_1,t_2}(x-y,z_1,z_2)|^2\Big(\frac{t_2}{t_2+|y_2|}\Big)^{n\lambda_1}\frac{dz_2dy_2}{t_2^n}\frac{dx_2dt_2}{t_2}\Big)^{1/2} \lesssim \frac{t_1^\alpha}{(t_1+|x_1-y_1-z_1|)^{n+\alpha}}.$$

(2) Combinations of Carleson and Hölder conditions :

$$\Big(\iint_{\widehat{I}}\int_{\mathbb{R}^n}\int_I |K_{t_1,t_2}(x-y,z_1,z_2)-K_{t_1,t_2}(x-y,z_1,z_2')|^2\Big(\frac{t_1}{t_1+|y_1|}\Big)^{n\lambda_1}\frac{dz_1dy_1}{t_1^n}\frac{dx_1dt_1}{t_1}\Big)^{1/2} \lesssim \frac{|z_2-z_2'|^\beta}{(t_2+|x_2-y_2-z_2|)^{m+\beta}},$$

whenever $|z_2-z_2'|<t_2/2$. And

$$\Big(\iint_{\widehat{J}}\int_{\mathbb{R}^m}\int_J |K_{t_1,t_2}(x-y,z_1,z_2)-K_{t_1,t_2}(x-y,z_1',z_2)|^2\Big(\frac{t_2}{t_2+|y_2|}\Big)^{m\lambda_2}\frac{dz_2dy_2}{t_2^n}\frac{dx_2dt_2}{t_2}\Big)^{1/2} \lesssim \frac{|z_1-z_1'|^\alpha}{(t_1+|x_1-y_1-z_1|)^{n+\alpha}},$$

whenever $|z_1-z_1'|<t_1/2$.

Assumption 6 (Bi-parameter Carleson condition of K_{t_1,t_2}). Let $\mathcal{D}=\mathcal{D}_n\times\mathcal{D}_m$, where $\mathcal{D}_n$ is a dyadic grid in $\mathbb{R}^n$ and $\mathcal{D}_m$ is a dyadic grid in $\mathbb{R}^m$. For $I\in\mathcal{D}_n$, let $W_I=I\times(\ell(I)/2,\ell(I))$ be the associated Whitney region. Denote $n_1=n, n_2=m$ and

$$C_{IJ}^{\mathcal{D}}=\iint_{W_J}\iint_{W_I}\iint_{\mathbb{R}^{n+m}}|\theta_{t_1,t_2}\mathbf{1}(y_1,y_2)|^2\Big[\prod_{i=1}^2\Big(\frac{t_i}{t_i+|x_i-y_i|}\Big)^{n_i\lambda_i}\Big]\frac{dy_1dy_2}{t_1^nt_2^m}\frac{dx_1dt_1}{t_1}\frac{dx_2dt_2}{t_2}.$$

We assume the following *bi-parameter Carleson condition*: For every $\mathcal{D}=\mathcal{D}_n\times\mathcal{D}_m$ there holds that

$$\sum_{\substack{I\times J\in\mathcal{D}\\ I\times J\subset\Omega}} C_{IJ}^{\mathcal{D}}\lesssim|\Omega| \tag{1.2}$$

for all sets $\Omega\subset\mathbb{R}^{n+m}$ such that $|\Omega|<\infty$ and such that for every $x\in\Omega$ there exists $I\times J\in\mathcal{D}$ so that $x\in I\times J\subset\Omega$.

The $L^2(\mathbb{R}^n\times\mathbb{R}^m)$ boundedness of bi-parameter g function is given by Martikainen [18], where the kernel s_{t_1,t_2} satisfies the **Assumptions 1**, **2**, **3**. Cao and Xue [1] establish the $L^2(\mathbb{R}^n\times\mathbb{R}^m)$ boundedness for bi-parameter $g_{\vec{\lambda}}^*$-function, where the kernel K_{t_1,t_2} satisfies the **Assumptions 4**, **5**, **6**. It deserves to note that the **Assumptions 3** is a necessary condition for the L^2 boundedness of bi-parameter g-function to be held, and **Assumptions 6** is also necessary for the L^2 boundedness of bi-parameter $g_{\vec{\lambda}}^*$-function.

Remark 1.5. It is easy to see that Assumptions 2 is a little weaker than Assumptions $2'$ and the same is true for Assumptions 5 and Assumption $5'$.

Now we state the main result of this paper.

Theorem 1.1. *Assume that the kernel s_{t_1,t_2} satisfies the **Assumptions 1**, **2′**, **3**. Then, it holds that*

$$\big\|g(f)\big\|_{L^1(\mathbb{R}^n\times\mathbb{R}^m)} \lesssim \big\|f\big\|_{H^1(\mathbb{R}^n\times\mathbb{R}^m)}, \tag{1.3}$$

where the implied constant depends only on the assumptions.

Theorem 1.2. *Let $\lambda_1,\lambda_2>2$, $0<\alpha\le n(\lambda_1-2)/2$ and $0<\beta\le m(\lambda_2-2)/2$. Assume that the kernel K_{t_1,t_2} satisfies the **Assumptions 4**, **5′**, **6**. Then, it holds that*

$$\big\|g^*_{\vec{\lambda}}(f)\big\|_{L^1(\mathbb{R}^n\times\mathbb{R}^m)} \lesssim \big\|f\big\|_{H^1(\mathbb{R}^n\times\mathbb{R}^m)}, \tag{1.4}$$

where the implied constant depends only on the assumptions.

By interpolation, we have the following corollaries:

Corollary 1.3. *Assume that the kernel s_{t_1,t_2} satisfies the **Assumptions 1**, **2′**, **3**. Then g is bounded on $L^p(\mathbb{R}^n\times\mathbb{R}^m)$ for all $1<p<2$.*

Corollary 1.4. *Let $\lambda_1,\lambda_2>2$, $0<\alpha\le n(\lambda_1-2)/2$ and $0<\beta\le m(\lambda_2-2)/2$. Assume that the kernel K_{t_1,t_2} satisfies the **Assumptions 4**, **5′**, **6**. Then $g^*_{\vec{\lambda}}$ is bounded on $L^p(\mathbb{R}^n\times\mathbb{R}^m)$ for all $1<p<2$.*

The organizations of this paper are as follows: Section 2 and Section 3 will be devoted to demonstrate Theorem 1.1 and Theorem 1.2. In section 4, we will briefly give the proofs of the Corollaries.

2. Proof of Theorem 1.1

In order to demonstrate Theorem 1.1, we first introduce some notions and a key lemma. Let $\mathcal{H}_i$ $(i=1,2)$ be the Hilbert space defined by

$$\mathcal{H}_i=\{f:(0,\infty)\times(0,\infty)\to\mathbb{C}\quad\text{is measurable},\quad |f|_{\mathcal{H}_i}<\infty\},$$

where

$$|f|_{\mathcal{H}_1}=\bigg\{\int_0^\infty\int_0^\infty |f(t_1,t_2)|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\bigg\}^{1/2}$$

and

$$|f|_{\mathcal{H}_2}=\bigg\{\int_{\mathbb{R}_+^{m+1}}\int_{\mathbb{R}_+^{n+1}} |f(t_1,t_2,y_1,y_2)|^2\frac{dy_1dt_1}{t_1}\frac{dy_2dt_2}{t_2}\bigg\}^{1/2}.$$

Hence, g and $g^*_{\vec{\lambda}}$ can be written by

$$g(f)(x)=|\{\theta_{t_1,t_2}f(x_1,x_2)\}_{t_1,t_2>0}|_{\mathcal{H}_1}:=|T_1|_{\mathcal{H}_1},$$

and

$$\begin{aligned} g^*_{\vec{\lambda}}(f)(x)&=\bigg|\bigg\{\frac{1}{t_1^nt_2^m}\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{\frac{n\lambda_1}{2}}\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{\frac{m\lambda_2}{2}}\vartheta_{t_1,t_2}f(x_1,x_2)\bigg\}_{t_1,t_2>0}\bigg|_{\mathcal{H}_2}\\ &:=|T_2|_{\mathcal{H}_2}. \end{aligned}$$

For $p>0$ and $i=1,2$, we define

$$L^p_{\mathcal{H}_i}(\mathbb{R}^n\times\mathbb{R}^m)=\left\{T_i(x_1,x_2):\mathbb{R}^n\times\mathbb{R}^m\to\mathcal{H}_i \ \text{ and } \ \|T_i\|_{L^p_{\mathcal{H}_i}(\mathbb{R}^n\times\mathbb{R}^m)}<\infty\right\}$$

where $\|T_i\|_{L^p_{\mathcal{H}_i}(\mathbb{R}^n\times\mathbb{R}^m)}=\left\{\iint_{\mathbb{R}^n\times\mathbb{R}^m}|T_i|^p_{\mathcal{H}_i}dx_1dx_2\right\}^{1/p}$.

The following vector-valued version of Theorem B provide a foundation for our proof.

Lemma 2.1. *Let T_i $(i=1,2)$ be an $\mathcal{H}_i$-valued linear operator which is bounded from $L^2(\mathbb{R}^n\times\mathbb{R}^m)$ to $L^2_{\mathcal{H}_i}(\mathbb{R}^n\times\mathbb{R}^m)$, and $0<p\le 1$. Suppose further that there exist constants $C>0$ such that for all $H^p(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atoms a supported on R, it holds that*

$$\iint_{\mathbb{R}^n\times\mathbb{R}^m\setminus\widetilde{R}_\gamma}|T_i(a)(x_1,x_2)|^p_{\mathcal{H}_i}dx_1dx_2\le C\gamma^{-\delta} \ \text{ for all } \ \gamma\ge 2 \ \text{ and some fixed } \ \delta>0,$$

where $\widetilde{R}_\gamma$ denotes the γ fold enlargement of R. Then there exists a constant $C>0$ such that for all $H^p(\mathbb{R}^n\times\mathbb{R}^m)$-atom $\widetilde{a}$, we have

$$\|T_i(\widetilde{a})\|_{L^p_{\mathcal{H}_i}(\mathbb{R}^n\times\mathbb{R}^m)}\le C.$$

Remark 2.1. The proof of Lemma 2.1 follows from the same idea of R. Fefferman. One only needs to replace $T(a)$ and $L^p(\mathbb{R}^n\times\mathbb{R}^m)$ in the proof given in [5] by $T_i(a)$ and $L^p_{\mathcal{H}_i}(\mathbb{R}^n\times\mathbb{R}^m)$, respectively. Here, we omit the proof.

Proof of Theorem 1.1. It suffices to verify that for any $H^1(\mathbb{R}^n\times\mathbb{R}^m)$-atom $\widetilde{a}$, there exists a constant $C>0$, such that

$$\|g(\widetilde{a})\|_{L^1(\mathbb{R}^n\times\mathbb{R}^m)}=\|T_1(\widetilde{a})\|_{L^1_{\mathcal{H}_1}(\mathbb{R}^n\times\mathbb{R}^m)}\le C. \tag{2.1}$$

In fact, for any $f\in H^1(\mathbb{R}^n\times\mathbb{R}^m)$, using atomic decomposition (Theorem A), we may split f by $f=\sum_j^\infty\lambda_j\widetilde{a}_j$, where $\widetilde{a}_j$ is $H^1(\mathbb{R}^n\times\mathbb{R}^m)$-atom. By the size condition of s_{t_1,t_2} and Fubini theorem, for all $t_1,t_2>0$, we have

$$\begin{aligned}
&\Big|\sum_{j=1}^\infty\lambda_j\iint_{\mathbb{R}^n\times\mathbb{R}^m}T_1(\widetilde{a}_j)dx_1dx_2\Big|\\
&\lesssim\sum_{j=1}^\infty|\lambda_j|\iint_{\mathbb{R}^n\times\mathbb{R}^m}\iint_{\mathbb{R}^n\times\mathbb{R}^m}|s_{t_1,t_2}(x_1,x_2,y_1,y_2)|dx_1dx_2|\widetilde{a}_j(y_1,y_2)|dy_1dy_2\\
&\lesssim\sum_{j=1}^\infty|\lambda_j|\iint_{\mathbb{R}^n\times\mathbb{R}^m}\iint_{\mathbb{R}^n\times\mathbb{R}^m}\frac{t_1^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{t_2^\beta}{(t_2+|x_2-y_2|)^{n+\beta}}dx_1dx_2\\
&\qquad|\widetilde{a}_j(y_1,y_2)|dy_1dy_2\\
&\lesssim\sum_{j=1}^\infty|\lambda_j|\iint_{\mathbb{R}^n\times\mathbb{R}^m}|\widetilde{a}_j(y_1,y_2)|dy_1dy_2\lesssim\sum_{j=1}^\infty|\lambda_j|.
\end{aligned}$$

Therefore, by Lebesgue dominated convergence theorem, it implies that

$$T_1(f)=\sum_{j=1}^\infty\lambda_jT_1(\widetilde{a}_j),\quad a.e. \tag{2.2}$$

Thus, by the above property (2.2), inequality (2.1) and Fatou's lemma, we have

$$\begin{aligned}\|T_1(f)\|_{L^1_{\mathcal{H}_1}(\mathbb{R}^n\times\mathbb{R}^m)} &= \int_{\mathbb{R}^n\times\mathbb{R}^m} |\sum_{j=1}^{\infty}\lambda_j T_1(\widetilde{a}_j)(x_1,x_2)|_{\mathcal{H}_1}dx_1dx_2\\ &\le \int_{\mathbb{R}^n\times\mathbb{R}^m}\sum_{j=1}^{\infty}|\lambda|_j|T_1(\widetilde{a}_j)(x_1,x_2)|_{\mathcal{H}_1}dx_1dx_2\\ &\le \sum_{j=1}^{\infty}|\lambda|_j\|T_1(\widetilde{a}_j)\|_{L^1_{\mathcal{H}_1}(\mathbb{R}^n\times\mathbb{R}^m)}\\ &\le \sum_{j=1}^{\infty}|\lambda|_j \lesssim \|f\|_{H^1(\mathbb{R}^n\times\mathbb{R}^m)}.\end{aligned} \tag{2.3}$$

Hence, to prove Theorem 1.1, it is sufficient to prove inequality (2.1).

Let a be a $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom, supported on a rectangle $R=I\times J$. Note that g-function is bounded on $L^2(\mathbb{R}^n\times\mathbb{R}^m)$. That is, T_1 is bounded from $L^2(\mathbb{R}^n\times\mathbb{R}^m)$ to $L^2_{\mathcal{H}_1}(\mathbb{R}^n\times\mathbb{R}^m)$. Thus, by Lemma 2.1, in order to prove inequality (2.1), it suffices to verify that

$$\iint_{\mathbb{R}^n\times\mathbb{R}^m\setminus\widetilde{R}_\gamma}|g(a)(x_1,x_2)|dx_1dx_2\lesssim\gamma^{-\delta},\qquad\text{for all }\gamma\ge 2. \tag{2.4}$$

Set $\gamma_1,\gamma_2\ge 2$ and $\gamma_1,\gamma_2\sim\gamma$, it's easy to see that

$$\begin{aligned}\widetilde{R}_\gamma \subset\ &\{(x_1,x_2)\in\mathbb{R}^n\times\mathbb{R}^m : x_1\notin\gamma_1 I, x_2\notin\gamma J\}\cup\{(x_1,x_2)\in\mathbb{R}^n\times\mathbb{R}^m : x_1\in\gamma_1 I, x_2\notin\gamma J\}\\ &\cup\{(x_1,x_2)\in\mathbb{R}^n\times\mathbb{R}^m : x_1\notin\gamma I, x_2\notin\gamma_2 J\}\cup\{(x_1,x_2)\in\mathbb{R}^n\times\mathbb{R}^m : x_1\notin\gamma I, x_2\in\gamma_2 J\}\}\\ &=: E_1\cup E_2\cup E_3\cup E_4.\end{aligned}$$

By symmetry, to prove (2.4), and thus to finish the proof of Theorem 1.1, we only need to prove that there exists a positive δ, such that

$$\iint_{E_1}|g(a)(x_1,x_2)|dx_1dx_2\lesssim\gamma^{-\delta}\ \text{ for all }\gamma\ge 2 \tag{2.5}$$

and

$$\iint_{E_2}|g(a)(x_1,x_2)|dx_1dx_2\lesssim\gamma^{-\delta}\ \text{ for all }\gamma\ge 2. \tag{2.6}$$

Now let us begin with the proof of inequality (2.5).

• **Proof of (2.5).** By the definition of g and support condition of a, we get

$$|g(a)(x)|^2=\int_0^\infty\int_0^\infty|\iint_{I\times J}s_{t_1,t_2}(x_1,x_2,y_1,y_2)a(y_1,y_2)dy_1\ dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}.$$

In order to estimate $|g(a)(x)|^2$ (where $x\in E_1$), we splitting the domain of variable t_1 and t_2 as follows,

$$\begin{aligned}(t_1,t_2)\in\mathbb{R}^+\times\mathbb{R}^+\subset&\big([|x_1-z_1|,\infty)\times[|x_2-z_2|,\infty)\big)\cup\big([|x_1-z_1|,\infty)\times(0,|x_2-z_2|)\big)\\ &\cup\big((0,|x_1-z_1|)\times[|x_2-z_2|,\infty)\big)\cup\big((0,|x_1-z_1|)\times(0,|x_2-z_2|)\big)\\ &=: F_1\cup F_2\cup F_3\cup F_4.\end{aligned}$$

Hence, we may write

$$|g(a)(x)|^2 \le \sum_{i=1}^{4}\int_0^\infty\int_0^\infty |\iint_{I\times J}\mathbf{1}_{F_i}(t_1,t_2)s_{t_1,t_2}(x_1,x_2,y_1,y_2)a(y_1,y_2)dy_1\ dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}$$
$$=:\sum_{i=1}^{4}A_i.$$

Let z_1 be the centre of I and z_2 be the centre of J, denote l_I and l_J as the sidelength of I and J, respectively. Note that, if $x\in E_1$ and $y\in I\times J$, then $|x_1-y_1|\sim|x_1-z_1|$ and $|x_2-y_2|\sim|x_2-z_2|$. We will continue to use the above notions in the rest of the paper, moreover, we always denote

$$\Phi^a_{t_1,t_2}(x,y,z):=\frac{|y_1-z_1|^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{|y_2-z_2|^\beta}{(t_2+|x_2-y_2|)^{m+\beta}}|a(y_1,y_2)|$$

Estimate for A_1. Since $x\in E_1$ and $(t_1,t_2)\in F_1$, then for any $y_1\in I$, we have $t_1\ge|x_1-z_1|\ge\gamma l_I\ge 2|y_1-z_1|$ and $t_1\gtrsim|x_1-y_1|$. Similarly, for any $y_2\in J$, we have $t_1\ge 2|y_1-z_1|$. The definition of A_1, vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom and Hölder estimate of s_{t_1,t_2} imply that

$$\begin{aligned}A_1&=\int_0^\infty\int_0^\infty|\iint_{I\times J}\mathbf{1}_{F_1}(t_1,t_2)(s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,z_2))\\&\quad-s_{t_1,t_2}(x,(z_1,y_2))+s_{t_1,t_2}(x,(z_1,z_2)))a(y_1,y_2)d\vec{y}|^2\frac{dt_1dt_2}{t_1t_2}\\&\lesssim\int_{|x_1-z_1|}^\infty\int_{|x_2-z_2|}^\infty|\iint_{I\times J}\Phi^a_{t_1,t_2}(x,y,z)dy_1dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\&\lesssim l_I^{2\alpha}l_J^{2\beta}\int_{|x_1-z_1|}^\infty\int_{|x_2-z_2|}^\infty\frac{dt_1}{t_1^{2n+2\alpha+1}}\frac{dt_2}{t_2^{2n+2\beta+1}}|\iint_{I\times J}|a(y_1,y_2)|dy_1dy_2|^2\\&\lesssim l_I^{2\alpha}l_J^{2\beta}|x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta}.\end{aligned}$$

Estimate for A_2. In this case, $x\in E_1$ and $(t_1,t_2)\in F_2$. Note that

$$\begin{aligned}F_2&\subset\big([|x_1-z_1|,\infty)\times(0,2|y_2-z_2|)\big)\cup\big([|x_1-z_1|,\infty)\times(2|y_2-z_2|,|x_2-z_2|)\big)\\&=:F_{2,1}\cup F_{2,2}.\end{aligned}$$

Then, one may deduce that

$$\begin{aligned}A_2&\le\sum_{i=1}^{2}\int_0^\infty\int_0^\infty|\iint_{I\times J}\mathbf{1}_{F_{2,i}}(t_1,t_2)s_{t_1,t_2}(x_1,x_2,y_1,y_2)a(y_1,y_2)dy_1\ dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\&=:A_{2,1}+A_{2,2}.\end{aligned}$$

First, we consider the contribution of $A_{2,1}$. Note that if $x\in E_1$ and $(t_1,t_2)\in F_{2,1}$, then for any $y_1\in I$, it holds that $t_1\ge 2|y_1-z_1|$ and $t_1\gtrsim|x_1-y_1|\sim|x_1-z_1|$. In addition, $t_2<2|y_2-z_2|<2l_J<|x_2-y_2|\sim|x_2-z_2|$. By the vanishing and size condition of

$H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom and mixed Hölder and size conditions of s_{t_1,t_2}, we obtain

$$\begin{aligned}
A_{2,1}&=\int_0^\infty\int_0^\infty|\iint_{I\times J}\mathbf{1}_{F_{2,1}}(t_1,t_2)(s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(z_1,y_2)))a(y_1,y_2)d\vec{y}|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\
&\lesssim\int_{|x_1-z_1|}^\infty\int_0^{2|y_2-z_2|}[\iint_{I\times J}\frac{|y_1-z_1|^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{t_2^\beta}{(t_2+|x_2-y_2|)^{m+\beta}}|a(y_1,y_2)|d\vec{y}]^2\frac{dt_1t_2}{t_1t_2}\\
&\lesssim l_I^{2\alpha}|x_2-z_2|^{-2m-2\beta}\int_{|x_1-z_1|}^\infty\int_0^{2|y_2-z_2|}\frac{t_2^{\beta-1}}{t_1^{2n+2\alpha+1}}dt_1dt_2|\iint_{I\times J}|a(y_1,y_2)|dy_1dy_2|^2\\
&\lesssim l_I^{2\alpha}l_J^{2\beta}|x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta}.
\end{aligned}$$

Nextly, we consider $A_{2,2}$.

Since $x\in E_1$ and $(t_1,t_2)\in F_{2,2}$, then for any $y_1\in I$, we have $t_1\geq|x_1-z_1|\geq\gamma l_I\geq 2|y_1-z_1|$ and $t_1\gtrsim|x_1-y_1|$. Similarly, for any $y_2\in J$ satisfies $|x_2-y_2|\geq t_1\geq 2|y_1-z_1|$. Set $\varepsilon<\min\{\alpha,\beta\}$. The vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom and Hölder estimate of s_{t_1,t_2} yield that

$$\begin{aligned}
A_{2,2}&=\int_0^\infty\int_0^\infty|\iint_{I\times J}\mathbf{1}_{F_{2,2}}(t_1,t_2)(s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,z_2))\\
&\quad-s_{t_1,t_2}(x,(z_1,y_2))+s_{t_1,t_2}(x,(z_1,z_2)))a(y_1,y_2)dy_1\ dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\
&\lesssim\int_{|x_1-z_1|}^\infty\int_{2|y_2-z_2|}^{|x_2-y_2|}|\iint_{I\times J}\Phi^a_{t_1,t_2}(x,y,z)dy_1dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\
&\lesssim l_I^{2\alpha}l_J^{2\beta}\big(\int_{|x_1-z_1|}^\infty\frac{dt_1}{t_1^{2n+2\alpha+1}}\big)\int_{2|y_2-z_2|}^{|x_2-y_2|}|\iint_{I\times J}\frac{|x_2-y_2|^\varepsilon}{t_2^\varepsilon|x_2-y_2|^{m+\beta}}|a(y_1,y_2)|dy_1dy_2|^2\frac{dt_2}{t_2}\\
&\lesssim l_I^{2\alpha}l_J^{2\beta-2\varepsilon}|x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta+2\varepsilon}.
\end{aligned}$$

Thus, it yields that

$$\begin{aligned}
A_2&\lesssim A_{2,1}+A_{2,2}\\
&\lesssim l_I^{2\alpha}l_J^{2\beta}|x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta}+l_I^{2\alpha}l_J^{2\beta-2\varepsilon}|x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta+2\varepsilon}.
\end{aligned}$$

Estimate for A_3. Since A_3 is symmetric with A_2, we may obtain that

$$A_3\lesssim l_I^{2\alpha}l_J^{2\beta}|x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta}+l_I^{2\alpha-2\varepsilon}l_J^{2\beta}|x_1-z_1|^{-2n-2\alpha+2\varepsilon}|x_2-z_2|^{-2m-2\beta}.$$

Estimate for A_4. Recall that $F_4=(0,|x_1-z_1|)\times(0,|x_2-z_2|)$. In order to estimate A_4, we further split F_4 as follows,

$$\begin{aligned}
F_4&\subset\big((2|y_1-z_1|,|x_1-z_1|)\times(2|y_1-z_1|,|x_2-z_2|)\big)\cup\big((0,2|y_1-z_1|)\times(0,2|y_2-z_2|)\big)\\
&\quad\cup\big((2|y_1-z_1|,|x_1-z_1|)\times(0,2|y_2-z_2|)\big)\cup\big((0,2|y_1-z_1|)\times(2|y_1-z_1|,|x_1-z_1|)\big)\\
&=:F_{4,1}\cup F_{4,2}\cup F_{4,3}\cup F_{4,4}.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}A_4 &\le \sum_{i=1}^{4}\int_0^\infty\int_0^\infty \Big|\iint_{I\times J}\mathbf{1}_{F_{4,i}}(t_1,t_2)s_{t_1,t_2}(x_1,x_2,y_1,y_2)a(y_1,y_2)dy_1dy_2\Big|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\ &=:\sum_{i=1}^{4}A_{4,i}.\end{aligned}$$

Now, we begin with the estimate of $A_{4,1}$.

Note that if $x\in E_1$ and $(t_1,t_2)\in F_{4,1}$, then for any $y_1\in I$ and $y_2\in J$, we have $|x_1-z_1|\ge t_1\ge 2|y_1-z_1|$ and $|x_2-z_2|\ge t_2\ge 2|y_2-z_2|$. Thus, by the vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom and Hölder conditions of s_{t_1,t_2}, we get

$$\begin{aligned}A_{4,1} &= \int_0^\infty\int_0^\infty \Big|\iint_{I\times J}\mathbf{1}_{F_{4,1}}(t_1,t_2)(s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,z_2))\\ &\quad -s_{t_1,t_2}(x,(z_1,y_2))+s_{t_1,t_2}(x,(z_1,z_2)))a(y_1,y_2)dy_1dy_2\Big|^2\frac{dt_1dt_2}{t_1t_2}\\ &\lesssim \int_{2|y_1-z_1|}^{|x_1-y_1|}\int_{2|y_2-z_2|}^{|x_2-y_2|}\Big|\iint_{I\times J}\Phi^a_{t_1,t_2}(x,y,z)dy_1dy_2\Big|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\ &\lesssim l_I^{2\alpha}l_J^{2\beta}\int_{2|y_1-z_1|}^{|x_1-y_1|}\int_{2|y_2-z_2|}^{|x_2-y_2|}\Big|\iint_{I\times J}\frac{|x_1-y_1|^\varepsilon}{t_1^\varepsilon|x_1-y_1|^{m+\beta}}\frac{|x_2-y_2|^\varepsilon}{t_2^\varepsilon|x_2-y_2|^{m+\beta}}\\ &\quad\times|a(y_1,y_2)|dy_1dy_2\Big|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\ &\lesssim l_I^{2\alpha-2\varepsilon}l_J^{2\beta-2\varepsilon}|x_1-z_1|^{-2n-2\alpha+2\varepsilon}|x_2-z_2|^{-2m-2\beta+2\varepsilon}.\end{aligned}$$

The vanishing and size conditions of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom and size estimate of s_{t_1,t_2} yield that

$$\begin{aligned}A_{4,2} &= \int_0^\infty\int_0^\infty \Big|\iint_{I\times J}\mathbf{1}_{F_{4,2}}(t_1,t_2)s_{t_1,t_2}(x,y)a(y_1,y_2)dy_1dy_2\Big|^2\frac{dt_1t_2}{t_1t_2}\\ &\lesssim \int_0^{2|y_1-z_1|}\int_0^{2|y_2-z_2|}\Big|\iint_{I\times J}\frac{t_1^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}}\frac{t_2^\beta}{(t_2+|x_2-y_2|)^{m+\beta}}\\ &\quad\times|a(y_1,y_2)|dy_1dy_2\Big|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\ &\lesssim |x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta}\int_{2|y_1-z_1|}^{|x_1-y_1|}\int_{2|y_2-z_2|}^{|x_2-y_2|}t_1^{-1-2\alpha}t_2^{-1-2\beta}dt_1dt_2\\ &\lesssim l_I^{2\alpha}l_J^{2\beta}|x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta}.\end{aligned}$$

Now, we are in the position to estimate $A_{4,3}$.

Note that if $x\in E_1$ and $(t_1,t_2)\in F_{4,3}$, then for any $y_1\in I$, it holds that $|x_1-z_1|\ge t_1\ge 2|y_1-z_1|$. By the vanishing and size conditions of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom

and Hölder conditions of s_{t_1,t_2}, we get

$$\begin{aligned}
A_{4,3} &= \int_0^\infty \int_0^\infty \Big| \iint_{I\times J} \mathbf{1}_{F_{4,3}}(t_1,t_2)(s_{t_1,t_2}(x,y) - s_{t_1,t_2}(x,(z_1,y_2)))a(y_1,y_2)dy_1dy_2\Big|^2 \frac{dt_1t_2}{t_1t_2} \\
&\lesssim \int_{2|y_1-z_1|}^{|x_1-y_1|} \int_0^{2|y_2-z_2|} \Big| \iint_{I\times J} \frac{|y_1-z_1|^\alpha}{(t_1+|x_1-y_1|)^{n+\alpha}} \frac{t_2^\beta}{(t_2+|x_2-y_2|)^{m+\beta}} \\
&\quad \times |a(y_1,y_2)|dy_1dy_2\Big|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \\
&\lesssim l_I^{2\alpha} l_J^{2\beta} |x_2-z_2|^{-2n-2\beta} \int_{2|y_1-z_1|}^{|x_1-y_1|} \Big| \iint_{I\times J} \frac{|x_1-y_1|^\varepsilon}{t_1^\varepsilon |x_1-y_1|^{n+\alpha}} |a(y_1,y_2)|dy_1dy_2\Big|^2 \frac{dt_1}{t_1} \\
&\lesssim l_I^{2\alpha-2\varepsilon} l_J^{2\beta} |x_1-z_1|^{-2n-2\alpha+2\varepsilon} |x_2-z_2|^{-2m-2\beta}.
\end{aligned}$$

By the fact that $A_{4,4}$ is symmetry with $A_{4,3}$, we may obtain

$$A_{4,4} \lesssim l_I^{2\alpha} l_J^{2\beta-2\varepsilon} |x_1-z_1|^{-2n-2\alpha} |x_2-z_2|^{-2m-2\beta+2\varepsilon}.$$

Therefore, we conclude that

$$\begin{aligned}
|g(a)(x)|^2 \le \sum_{i=1}^4 A_i \lesssim\ & l_I^{2\alpha-2\varepsilon} l_J^{2\beta-2\varepsilon} |x_1-z_1|^{-2n-2\alpha+2\varepsilon} |x_2-z_2|^{-2m-2\beta+2\varepsilon} \\
&+ l_I^{2\alpha} l_J^{2\beta} |x_1-z_1|^{-2n-2\alpha} |x_2-z_2|^{-2m-2\beta} \\
&+ l_I^{2\alpha-2\varepsilon} l_J^{2\beta} |x_1-z_1|^{-2n-2\alpha+2\varepsilon} |x_2-z_2|^{-2m-2\beta} \\
&+ l_I^{2\alpha} l_J^{2\beta-2\varepsilon} |x_1-z_1|^{-2n-2\alpha} |x_2-z_2|^{-2m-2\beta+2\varepsilon}.
\end{aligned}$$

Recall that $\varepsilon < \min\{\alpha,\beta\}$ and $\gamma_1 \sim \gamma$. The above estimate leads to

$$\iint_{E_1} |g(a)(x_1,x_2)|dx_1dx_2 \lesssim \gamma^{-\min\{\alpha,\beta\}}.$$

• **Proof of (2.6).** Recall that $E_2 = \{(x_1,x_2) \in \mathbb{R}^n \times \mathbb{R}^m : x_1 \in \gamma_1 I, x_2 \notin \gamma J\}$, we have

$$\begin{aligned}
E_2 \subset\ & \{(x_1,x_2) \in \mathbb{R}^n \times \mathbb{R}^m : x_1 \in 2I, x_2 \notin \gamma J\} \\
& \cup \{(x_1,x_2) \in \mathbb{R}^n \times \mathbb{R}^m : x_1 \in \gamma_1 I \backslash 2I, x_2 \notin \gamma J\} =: E_{2,1} \cup E_{2,2}.
\end{aligned}$$

Therefore, it holds that

$$\iint_{E_2} |g(a)(x_1,x_2)|dx_1dx_2 \lesssim \iint_{E_{2,1}} |g(a)(x_1,x_2)|dx_1dx_2 + \iint_{E_{2,2}} |g(a)(x_1,x_2)|dx_1dx_2.$$

Since $E_{2,2} \subset \{(x_1,x_2) \in \mathbb{R}^n \times \mathbb{R}^m : x_1 \notin 2I, x_2 \notin \gamma J\}$, taking $\gamma_1 = 2$ and repeating the proof of (2.5), we may get

$$\iint_{E_{2,2}} |g(a)(x_1,x_2)|dx_1dx_2 \lesssim \gamma^{\min\{\alpha,\beta\}}.$$

Hence, to complete the proof of Theorem 1.3, we only need to show that there exists a $\delta > 0$ such that

$$\iint_{E_{2,1}} |g(a)(x_1,x_2)|dx_1dx_2 \lesssim \gamma^{-\delta}. \tag{2.7}$$

To do this, similar to the analysis in the proof of (2.5), we split the domain of variable t_1 and t_2 as follows,

$$\begin{aligned}(t_1,t_2)\in\mathbb{R}^+\times\mathbb{R}^+ \subset&\big((0,2l_I]\times[|x_2-z_2|,\infty)\big)\cup\big((0,2l_I]\times(2|y_2-z_2|,|x_2-z_2|)\big)\\ &\cup\big((0,2l_I]\times[0,2|y_2-z_2|)\big)\cup\big((2l_I,\infty)\times[|x_2-z_2|,\infty)\big)\\ &\cup\big((2l_I,\infty)\times(2|y_2-z_2|,|x_2-z_2|)\big)\cup\big((2l_I,\infty)\times[0,2|y_2-z_2|)\big)\\ &=: G_1\cup G_2\cup G_3\cup G_4\cup G_5\cup G_6.\end{aligned}$$

Thus, we may obtain

$$\begin{aligned}\iint_{E_{2,1}}|g(a)(x_1,x_2)|dx_1dx_2\le\sum_{i=1}^6\iint_{E_2}\Big(\int_0^\infty\int_0^\infty\Big|\iint_{I\times J}\mathbf{1}_{G_i}(t_1,t_2)s_{t_1,t_2}(x_1,x_2,y_1,y_2)\\ \times a(y_1,y_2)dy_1\,dy_2\Big|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\Big)^{1/2}dx_1dx_2=:\sum_{i=1}^6 B_i.\end{aligned}$$

Now, let us begin with the estimate of B_1.

Estimate for B_1. Since $x\in E_2$ and $(t_1,t_2)\in G_1$, then for any $y_2\in J$, it satisfies that $t_2\ge 2|y_2-z_2|$. The support, vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom, Hölder inequality and the combinations of Carleson and Hölder conditions of s_{t_1,t_2} imply that

$$\begin{aligned}B_1=&\iint_{2I\times(\gamma J)^c}\bigg(\int_0^\infty\int_0^\infty\Big|\iint_{2I\times J}\mathbf{1}_{G_1}(t_1,t_2)(s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,z_2)))\\ &\times a(y_1,y_2)dy_1dy_2\Big|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\bigg)^{1/2}dx_1dx_2\\ \lesssim&\|a\|_{L^2}\iint_{2I\times(\gamma J)^c}\bigg(\int_0^{2l_I}\int_{|x_2-z_2|}^\infty\iint_{2I\times J}|s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,z_2))|^2\\ &\times dy_1dy_2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\bigg)^{1/2}dx_1dx_2\\ \lesssim&\|a\|_{L^2}(2I)^{1/2}\int_{(\gamma J)^c}\bigg(\int_{|x_2-z_2|}^\infty\int_J\Big(\iint_{\widehat{2I}}\int_{2I}|s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,z_2))|^2\\ &\times dy_1\frac{dx_1dt_1}{t_1}\Big)dy_2\frac{dt_2}{t_2}\bigg)^{1/2}dx_2\\ \lesssim&\|a\|_{L^2}I^{1/2}\int_{(\gamma J)^c}\bigg(\int_{|x_2-z_2|}^\infty\int_J\frac{|y_2-z_2|^{2\beta}}{(t_2+|x_2-y_2|)^{2m+2\beta}}dy_2\frac{dt_2}{t_2}\bigg)^{1/2}dx_2\\ \lesssim&\|a\|_{L^2}I^{1/2}J^{1/2}l_J^\beta\int_{(\gamma J)^c}|x_2-z_2|^{-m-\beta}dx_2\lesssim\gamma^{-\beta}.\end{aligned}\tag{2.8}$$

Now, we consider the estimate of B_2.

Estimate for B_2. If $x\in E_2$ and $(t_1,t_2)\in G_1$, then for any $y_2\in J$, it holds that $|x_2-z_2|\ge t_2\ge 2|y_2-z_2|$. Recall $\varepsilon\le\min\{\alpha,\beta\}$. Similar to inequality (2.8), by the support, vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom, Hölder inequality

and the combinations of Carleson and Hölder conditions of s_{t_1,t_2}, we may obtain

$$\begin{aligned}
B_2 &= \iint_{2I\times(\gamma J)^c}\Big(\int_0^\infty\int_0^\infty|\iint_{2I\times J}\mathbf{1}_{G_2}(t_1,t_2)(s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,z_2)))\\
&\qquad\times a(y_1,y_2)dy_1dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\Big)^{1/2}dx_1dx_2\\
&\lesssim \|a\|_{L^2}I^{1/2}\int_{(\gamma J)^c}\Big(\int_{2|y_2-z_2|}^{|x_2-z_2|}\int_J\frac{|y_2-z_2|^{2\beta}}{(t_2+|x_2-y_2|)^{2m+2\beta}}dy_2\frac{dt_2}{t_2}\Big)^{1/2}dx_2\\
&\lesssim \|a\|_{L^2}I^{1/2}l_J^\beta\int_{(\gamma J)^c}\Big(\int_{2|y_2-z_2|}^{|x_2-z_2|}\int_J\frac{|x_2-z_2|^{\varepsilon}}{t_2^{\varepsilon}|x_2-z_2|^{2m+2\beta}}dy_2\frac{dt_2}{t_2}\Big)^{1/2}dx_2\\
&\lesssim \|a\|_{L^2}I^{1/2}J^{1/2}l_J^{\beta-\varepsilon/2}\int_{(\gamma J)^c}|x_2-z_2|^{-m-\beta+\varepsilon/2}dx_2\lesssim\gamma^{-\beta/2}.
\end{aligned}$$

Estimate for B_3. The support, vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom, Hölder inequality and the Combinations of Carleson and size conditions of s_{t_1,t_2} yield that

$$\begin{aligned}
B_3 &= \iint_{2I\times(\gamma J)^c}\Big(\int_0^\infty\int_0^\infty|\iint_{2I\times J}\mathbf{1}_{G_3}(t_1,t_2)s_{t_1,t_2}(x,y)\\
&\qquad\times(y_1,y_2)dy_1dy_2|^2\frac{dt_1dt_2}{t_1t_2}\Big)^{1/2}dx_1dx_2\\
&\lesssim \|a\|_{L^2}(2I)^{1/2}\int_{(\gamma J)^c}\Big(\int_0^{2|y_2-z_2|}\int_J\big(\iint_{\widehat{2I}}\int_{2I}|s_{t_1,t_2}(x_1,x_2,y_1,y_2)|dy_1\frac{dx_1dt_1}{t_1}\big)\\
&\qquad\times dy_2\frac{dt_2}{t_2}\Big)^{1/2}dx_2\\
&\lesssim \|a\|_{L^2}I^{1/2}\int_{(\gamma J)^c}\Big(\int_0^{2|y_2-z_2|}\int_J\frac{t_2^{2\beta}}{(t_2+|x_2-y_2|)^{2m+2\beta}}dy_2\frac{dt_2}{t_2}\Big)^{1/2}dx_2\\
&\lesssim \|a\|_{L^2}I^{1/2}J^{1/2}l_J^{\beta}\int_{(\gamma J)^c}|x_2-z_2|^{-m-\beta}dx_2\lesssim\gamma^{-\beta}.
\end{aligned}$$

Estimate for B_4. Since $x\in E_2$ and $(t_1,t_2)\in G_4$, then for any $y_1\in I$, it satisfies that $t_1\geq 2|y_1-z_1|$ and $t_1>|x_1-y_1|$. For any $y_2\in J$, it satisfies that $t_2\geq|x_2-z_2|\geq 2|y_2-z_2|$. The vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom and Hölder estimate of s_{t_1,t_2} yield that

$$\begin{aligned}
&\int_0^\infty\int_0^\infty|\iint_{I\times J}\mathbf{1}_{G_4}(t_1,t_2)s_{t_1,t_2}(x,y))a(y_1,y_2)dy_1\ dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}\\
&=\int_0^\infty\int_0^\infty|\iint_{I\times J}\mathbf{1}_{G_4}(t_1,t_2)(s_{t_1,t_2}(x,y)-s_{t_1,t_2}(x,(y_1,z_2))\\
&\qquad-s_{t_1,t_2}(x,(z_1,y_2))+s_{t_1,t_2}(x,(z_1,z_2)))a(y_1,y_2)dy_1\ dy_2|^2\frac{dt_1}{t_1}\frac{dt_2}{t_2}
\end{aligned}\tag{2.9}$$

$$\begin{aligned}
&\lesssim \int_{2l_I}^{\infty}\int_{|x_2-z_2|}^{\infty} \Big| \iint_{I\times J} \frac{|y_1-z_1|^{\alpha}}{(t_1+|x_1-y_1|)^{n+\alpha}} \frac{|y_2-z_2|^{\beta}}{(t_2+|x_2-y_2|)^{m+\beta}} \\
&\quad \times |a(y_1,y_2)| dy_1 dy_2|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \\
&\lesssim l_I^{2\alpha} l_J^{2\beta} \int_{2l_I}^{\infty}\int_{|x_2-z_2|}^{\infty} \frac{dt_1}{t_1^{2n+2\alpha+1}} \frac{dt_2}{t_2^{2n+2\beta+1}} \Big| \iint_{I\times J} |a(y_1,y_2)| dy_1 dy_2|^2 \\
&\lesssim l_I^{-2n} l_J^{2\beta} |x_2-z_2|^{-2m-2\beta}.
\end{aligned}$$

Putting the above estimates into the definition of B_4, it holds that

$$B_4 \lesssim l_I^{-n} l_J^{\beta} \int_{2I}\int_{(\gamma J)^c} |x_2-z_2|^{-m-\beta} dx_1 dx_2 \lesssim \gamma^{-\beta}.$$

Estimate for B_5. Note that $x \in E_2$ and $(t_1,t_2) \in G_5$, then for any $y_1 \in I$, it satisfies that $t_1 \geq 2|y_1-z_1|$ and $t_1 > |x_1-y_1|$. For any $y_2 \in J$, it satisfies $|x_2-z_2| \geq t_2 \geq 2|y_2-z_2|$. Similar to inequality (2.9), by the vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom and Hölder estimate of s_{t_1,t_2}, we may get

$$\begin{aligned}
&\int_0^{\infty}\int_0^{\infty} \Big| \iint_{I\times J} \mathbf{1}_{G_5}(t_1,t_2) s_{t_1,t_2}(x,y)) a(y_1,y_2) dy_1\ dy_2|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \\
&\lesssim \int_{2l_I}^{\infty}\int_{2|y_2-z_2|}^{|x_2-z_2|} \Big| \iint_{I\times J} \frac{|y_1-z_1|^{\alpha}}{(t_1+|x_1-y_1|)^{n+\alpha}} \frac{|y_2-z_2|^{\beta}}{(t_2+|x_2-y_2|)^{m+\beta}} |a(y_1,y_2)| dy_1 dy_2|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \\
&\lesssim l_I^{2\alpha} l_J^{2\beta} \Big(\int_{2l_I}^{\infty} \frac{dt_1}{t_1^{2n+2\alpha+1}} \Big) \int_{2|y_2-z_2|}^{|x_2-y_2|} \Big| \iint_{I\times J} \frac{|x_2-y_2|^{\varepsilon/2}}{t_2^{\varepsilon/2}|x_2-y_2|^{m+\beta}} |a(y_1,y_2)| dy_1 dy_2|^2 \frac{dt_2}{t_2} \\
&\lesssim l_I^{-2n} l_J^{2\beta-\varepsilon} |x_2-z_2|^{-2m-2\beta+\varepsilon}.
\end{aligned}$$

This leads to

$$B_5 \lesssim l_I^{-n} l_J^{\beta-\varepsilon/2} \int_{2I}\int_{(\gamma J)^c} |x_2-z_2|^{-m-\beta+\varepsilon/2} dx_1 dx_2 \lesssim \gamma^{-\beta/2}.$$

Estimate for B_6. Note that $x \in E_2$ and $(t_1,t_2) \in G_6$, then for any $y_1 \in I$, it holds that $t_1 \geq 2|y_1-z_1|$ and $t_1 > |x_1-y_1|$. By the support, vanishing and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom and Hölder estimate of s_{t_1,t_2}, we may get

$$\begin{aligned}
&\int_0^{\infty}\int_0^{\infty} \Big| \iint_{I\times J} \mathbf{1}_{G_6}(t_1,t_2) s_{t_1,t_2}(x,y)) a(y_1,y_2) dy_1\ dy_2|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \\
&= \int_0^{\infty}\int_0^{\infty} \Big| \iint_{I\times J} \mathbf{1}_{F_{2,1}}(t_1,t_2) (s_{t_1,t_2}(x,y) - s_{t_1,t_2}(x,(z_1,y_2))) a(y_1,y_2) dy_1\ dy_2|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \\
&\lesssim \int_{2l_I}^{\infty}\int_0^{2|y_2-z_2|} \Big| \iint_{I\times J} \frac{|y_1-z_1|^{\alpha}}{(t_1+|x_1-y_1|)^{n+\alpha}} \frac{t_2^{\beta}}{(t_2+|x_2-y_2|)^{m+\beta}} |a(y_1,y_2)| dy_1 dy_2|^2 \frac{dt_1}{t_1}\frac{dt_2}{t_2} \\
&\lesssim l_I^{2\alpha} |x_2-z_2|^{-2m-2\beta} \int_{2l_I}^{\infty}\int_0^{2|y_2-z_2|} \frac{t_2^{\beta}}{t_1^{2n+2\alpha+1}} dt_1 dt_2 \Big| \iint_{I\times J} |a(y_1,y_2)| dy_1 dy_2|^2 \\
&\lesssim l_I^{2n} l_J^{2\beta} |x_2-z_2|^{-2m-2\beta}.
\end{aligned}$$

This yields that

$$B_6 \lesssim l_I^n l_J^\beta \int_{2I}\int_{(\gamma J)^c} |x_2 - z_2|^{-m-\beta} dx_1 dx_2 \lesssim \gamma^{-\beta}.$$

Combining the estimate for $B_i (i = 1 \cdots 6)$, we get

$$\iint_{E_{2,1}} |g(a)(x_1, x_2)| dx_1 dx_2 \lesssim \gamma^{-\beta/2}.$$

Thus, we finish the proof of the inequality (2.7). Since we have reduced the proof of Theorem 1.1 to inequality (2.7), the proof of Theorem 1.1 is also completed. □

3. Proof of Theorem 1.2

First, we list an elementary inequality which is useful in the proof of Theorem 1.2.

Lemma 3.1. ([19]) *Let $\theta_1, \theta_2 > 0$, $A \le B$. Denote $\theta_0 = \min\{\theta_1, \theta_2\}$, $\theta_3 = \max\{\theta_1, \theta_2\} > n$, then*

$$\int_{\mathbb{R}^n} \frac{dy}{(A + |y|)^{\theta_1}(B + |x - y|)^{\theta_2}} \lesssim A^{n-\theta_3}(|x| + B)^{-\theta_0}.$$

Proof of Theorem 1.2. Similar to Theorem 1.1, in order to prove (1.4), it suffices to verify that for any $H^1(\mathbb{R}^n \times \mathbb{R}^m)$-atom $\widetilde{a}$, there exists a constant $C > 0$ satisfying

$$\|g_\lambda^*(\widetilde{a})\|_{L^1(\mathbb{R}^n\times\mathbb{R}^m)} = \|T_2(\widetilde{a})\|_{L^1_{\mathcal{H}_2}(\mathbb{R}^n\times\mathbb{R}^m)} \le C. \tag{3.1}$$

In fact, let $f \in H^1(\mathbb{R}^n \times \mathbb{R}^m)$, by atomic decomposition, we have $f = \sum_j^\infty \lambda_j \widetilde{a}_j$, where $\widetilde{a}_j$ is $H^1(\mathbb{R}^n \times \mathbb{R}^m)$-atom. The size condition of K_{t_1,t_2} and Fubini theorem lead to

$$\begin{aligned}
&\sum_{j=1}^\infty \lambda_j \iint_{\mathbb{R}^n\times\mathbb{R}^m} T_2(\widetilde{a}_j)(x_1, x_2) dx_1 dx_2 \\
&\le \sum_{j=1}^\infty \lambda_j \frac{1}{t_1^n t_2^m} \iint_{\mathbb{R}^n\times\mathbb{R}^m}\iint_{\mathbb{R}^n\times\mathbb{R}^m} \Big(\frac{t_1}{t_1 + |x_1 - y_1|}\Big)^{\frac{n\lambda_1}{2}} \Big(\frac{t_2}{t_2 + |x_2 - y_2|}\Big)^{\frac{m\lambda_2}{2}} \\
&\qquad \times \iint_{\mathbb{R}^n\times\mathbb{R}^m} |K_{t_1,t_2}(y_1, y_2, z_1, z_2)| dy_1 dy_2 dx_1 dx_2 \widetilde{a}(z_1, z_2) dz_1 dz_2 \\
&\le \sum_{j=1}^\infty \frac{\lambda_j}{t_1^n t_2^m} \iint_{\mathbb{R}^n\times\mathbb{R}^m}\iint_{\mathbb{R}^n\times\mathbb{R}^m} \Big(\frac{t_1}{t_1 + |x_1 - y_1|}\Big)^{\frac{n\lambda_1}{2}} \Big(\frac{t_2}{t_2 + |x_2 - y_2|}\Big)^{\frac{m\lambda_2}{2}} \iint_{\mathbb{R}^n\times\mathbb{R}^m} \\
&\qquad \times \frac{t_1^\alpha}{(t_1 + |y_1 - z_1|)^{n+\alpha}} \frac{t_2^\beta}{(t_2 + |y_2 - z_2|)^{m+\beta}} dy_1 dy_2 dx_1 dx_2 |\widetilde{a}(z_1, z_2)| dz_1 dz_2.
\end{aligned} \tag{3.2}$$

Since $0 < \alpha \le n(\lambda_1 - 2)/2$ and $0 < \beta \le m(\lambda_2 - 2)/2$, we may set $\varepsilon_1 > 0$, $\varepsilon_2 > 0$ with $2n + 2\alpha \le 2n + 2\varepsilon_1 \le n\lambda_1$ and $2m + 2\beta \le 2m + 2\varepsilon_2 \le m\lambda_2$, respectively. By Lemma

3.1, we obtain

$$\begin{aligned}(3.3)\qquad &\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{\frac{n\lambda_1}{2}}\frac{t_1^\alpha}{(t_1+|y_1-z_1|)^{n+\alpha}}dy_1dx_1\\ &\lesssim\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{t_1^{n+\varepsilon_1}}{\big(t_1+|x_1-y_1|\big)^{n+\varepsilon_1}}\frac{t_1^\alpha}{(t_1+|y_1-z_1|)^{n+\alpha}}dy_1dx_1\\ &\lesssim\int_{\mathbb{R}^n}\frac{t_1^{n+\alpha}}{(t_1+|x_1-z_1|)^{n+\alpha}}dx_1\lesssim t_1^n.\end{aligned}$$

Similarly, we have

$$(3.4)\qquad \int_{\mathbb{R}^m}\int_{\mathbb{R}^m}\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{\frac{m\lambda_2}{2}}\frac{t_2^\beta}{(t_2+|y_2-z_2|)^{m+\beta}}dy_2dx_2\lesssim t_2^m.$$

Putting (3.3) and (3.4) into (3.5), we have

$$(3.5)\qquad \sum_{j=1}^\infty\lambda_j\iint_{\mathbb{R}^n\times\mathbb{R}^m}T_2(\widetilde{a}_j)(x_1,x_2)dx_1dx_2\le C.$$

Hence, (3.5) and Lebesgue dominated convergence theorem imply that

$$T_2(f)=\sum_{j=1}^\infty\lambda_jT_2(\widetilde{a}_j),\quad a.e.$$

By (3.1) and repeating the same steps as in (2.3), we have

$$\|T_2(f)\|_{L^1_{\mathcal{H}_2}(\mathbb{R}^n\times\mathbb{R}^m)}\lesssim\|f\|_{H^1(\mathbb{R}^n\times\mathbb{R}^m)}.$$

Thus, to prove Theorem 1.2, we only need to prove inequality (3.1).

From now on, we are devoted to prove (3.1).

Let a be any $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom, with support on a rectangle $R=I\times J$. As remarked before, $g^*_{\vec{\lambda}}(f)$ is bounded from $L^2(\mathbb{R}^n\times\mathbb{R}^m)$ to $L^2_{\mathcal{H}_2}(\mathbb{R}^n\times\mathbb{R}^m)$. Thus, by Lemma 2.1, in order to prove (1.4), it suffices to verify that

$$(3.6)\qquad \iint_{\mathbb{R}^n\times\mathbb{R}^m\setminus\widetilde{R}_\gamma}|g^*_{\vec{\lambda}}(a)(x_1,x_2)|dx_1dx_2\lesssim\gamma^{-\delta},\qquad\text{for all }\gamma\ge2.$$

Recall the definition of E_i in the proof of Theorem 1.1 and by symmetry, to prove (3.6), we only need to show that there exist a $\delta>0$ such that

$$(3.7)\qquad \iint_{E_1}|g^*_{\vec{\lambda}}(a)(x_1,x_2)|dx_1dx_2\lesssim\gamma^{-\delta}\ \text{ for all }\gamma\ge2$$

and

$$(3.8)\qquad \iint_{E_2}|g^*_{\vec{\lambda}}(a)(x_1,x_2)|dx_1dx_2\lesssim\gamma^{-\delta}\ \text{ for all }\gamma\ge2.$$

Now, let us to begin with the proof of (3.7).

• **Proof of (3.7).** By the definition of $g^*_{\vec{\lambda}}$ and the support condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$

rectangle atom, we get

$$|g^*_{\vec{\lambda}}(a)(x)|^2 = \iint_{\mathbb{R}^{m+1}_+} \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \iint_{\mathbb{R}^{n+1}_+} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \times |\iint_{\mathbb{R}^n\times\mathbb{R}^m} K_{t_1,t_2}(y_1,y_2,z_1,z_2)a(z_1,z_2)dz_1\ dz_2|^2 \frac{dy_1dt_1}{t_1^{n+1}}\frac{dy_2dt_2}{t_2^{m+1}}.$$

With abuse of notation, we split the domain of variable t_1 and t_2 as follows,

$$\begin{aligned}(t_1,t_2)\in\mathbb{R}^+\times\mathbb{R}^+ \subset &\big([2l_I,\infty)\times[2l_J,\infty)\big)\cup\big([2l_I,\infty)\times(0,2l_J)\big)\\ &\cup\big((0,2l_I)\times[2l_J,\infty)\big)\cup\big((0,2l_I)\times(0,2l_J)\big)\\ &=: F_1\cup F_2\cup F_3\cup F_4.\end{aligned}$$

Hence, for any $x\in E_1$, we have

$$\begin{aligned}|g^*_{\vec{\lambda}}(a)(x)|^2 &\le \sum_{i=1}^4 \iint_{\mathbb{R}^{m+1}_+} \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \iint_{\mathbb{R}^{n+1}_+} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \mathbf{1}_{F_i}(t_1,t_2)\\ &\quad |\iint_{\mathbb{R}^n\times\mathbb{R}^m} K_{t_1,t_2}(y_1,y_2,z_1,z_2)a(z_1,z_2)dz_1\ dz_2|^2 \frac{dy_1dt_1}{t_1^{n+1}}\frac{dy_2dt_2}{t_2^{m+1}}\\ &=: \sum_{i=1}^4 A_i.\end{aligned}$$

Denote z_1' as the centre of I and z_2' as the centre of J. Now, we will estimate each A_i. **Estimate for A_1.** Since $x\in E_1$ and $(t_1,t_2)\in F_1$, then for any $y_1\in I$, $y_2\in J$, we have $t_1\ge 2l_I\ge 2|z_1-z_1'|$ and $t_2\ge 2l_J\ge 2|z_2-z_2'|$. By the vanishing, support and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom, Hölder estimate of K_{t_1,t_2} and Hölder inequality, we have

$$\begin{aligned}A_1 &= \iint_{\mathbb{R}^{m+1}_+} \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \iint_{\mathbb{R}^{n+1}_+} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \mathbf{1}_{F_1}(t_1,t_2)\\ &\quad \times |\iint_{I\times J} (K_{t_1,t_2}(y,z)-K_{t_1,t_2}(y,(z_1,z_2'))\\ &\quad - K_{t_1,t_2}(y,(z_1',z_2))+K_{t_1,t_2}(y,(z_1',z_2')))a(z_1,z_2)dz_1dz_2|^2 \frac{dy_1dt_1}{t_1^{n+1}}\frac{dy_2dt_2}{t_2^{m+1}}\\ &\lesssim \int_{2l_J}^\infty\int_{\mathbb{R}^m} \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \int_{2l_I}^\infty\int_{\mathbb{R}^n} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} |\iint_{I\times J}\\ &\quad \frac{|z_1-z_1'|^\alpha}{(t_1+|y_1-z_1|)^{n+\alpha}}\frac{|z_2-z_2'|^\beta}{(t_2+|y_2-z_2|)^{m+\beta}}a(z_1,z_2)dz_1\ dz_2|^2 \frac{dy_1dt_1}{t_1^{n+1}}\frac{dy_2dt_2}{t_2^{m+1}}\\ &\lesssim \|a\|_2^2 \int_{2l_I}^\infty\int_{\mathbb{R}^n}\int_I \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \frac{|z_1-z_1'|^{2\alpha}}{(t_1+|y_1-z_1|)^{2n+2\alpha}}dz_1\frac{dy_1dt_1}{t_1^{n+1}}\\ &\quad \times \int_{2l_J}^\infty\int_{\mathbb{R}^m}\int_J \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \frac{|z_2-z_2'|^{2\beta}}{(t_2+|y_2-z_2|)^{2m+2\beta}}dz_2\frac{dy_2dt_2}{t_2^{m+1}}.\end{aligned} \tag{3.9}$$

By Lemma 3.1, it holds that

$$\begin{aligned}(3.10)\qquad &\int_{2l_I}^{\infty}\int_{\mathbb{R}^n}\int_I \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}\frac{|z_1-z_1'|^{2\alpha}}{(t_1+|y_1-z_1|)^{2n+2\alpha}}dz_1\frac{dy_1dt_1}{t_1^{n+1}}\\ &\lesssim l_I^{2\alpha}\int_{2l_I}^{\infty}\int_I\int_{\mathbb{R}^n}\frac{t_1^{2n+2\varepsilon_1}}{\big(t_1+|x_1-y_1|\big)^{2n+2\varepsilon_1}}\frac{1}{(t_1+|y_1-z_1|)^{2n+2\alpha}}dy_1\frac{dz_1dt_1}{t_1^{n+1}}\\ &\lesssim l_I^{2\alpha}\int_{2l_I}^{\infty}\int_I\frac{t_1^n}{\big(t_1+|x_1-z_1|\big)^{2n+2\alpha}}\frac{dz_1dt_1}{t_1^{n+1}}\lesssim l_I^{n+2\alpha}|x_1-z_1|^{-2n-2\alpha}.\end{aligned}$$

Similarly, we may get

$$\begin{aligned}(3.11)\qquad &\int_{2l_J}^{\infty}\int_{\mathbb{R}^m}\int_J\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\frac{|z_2-z_2'|^{2\beta}}{(t_2+|y_2-z_2|)^{2n+2\beta}}dz_2\frac{dy_2dt_2}{t_2^{m+1}}\\ &\qquad\lesssim l_J^{m+2\beta}|x_2-z_2|^{-2m-2\beta}.\end{aligned}$$

Putting the above estimate into (3.9) and recall that $\|a\|_2^2\lesssim|R|^{-1}=l_I^{-n}l_J^{-m}$, we have

$$A_1\lesssim l_I^{2\alpha}l_J^{2\beta}|x_1-z_1|^{-2n-2\alpha}|x_2-z_2|^{-2m-2\beta}.$$

Estimate for A_2. Since $x\in E_1$ and $(t_1,t_2)\in F_2$, then for any $y_1\in I$, we have $t_1\geq 2l_I\geq 2|z_1-z_1'|$. By the vanishing, support and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom, mixed Hölder and size condition of K_{t_1,t_2} and Hölder inequality, we have

$$\begin{aligned}(3.12)\qquad A_2=&\iint_{\mathbb{R}_+^{m+1}}\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\iint_{\mathbb{R}_+^{n+1}}\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}\mathbf{1}_{F_2}(t_1,t_2)\\ &|\iint_{I\times J}(K_{t_1,t_2}(y,z)-K_{t_1,t_2}(y,(z_1',z_2)))a(z_1,z_2)dz_1\ dz_2|^2\frac{dy_1dt_1}{t_1^{n+1}}\frac{dy_2dt_2}{t_2^{m+1}}\\ \lesssim&\int_0^{2l_J}\int_{\mathbb{R}^m}\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\int_{2l_I}^{\infty}\int_{\mathbb{R}^n}\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}|\iint_{I\times J}\\ &\frac{|z_1-z_1'|^{\alpha}}{(t_1+|y_1-z_1|)^{n+\alpha}}\frac{t_2^{\beta}}{(t_2+|y_2-z_2|)^{m+\beta}}a(z_1,z_2)dz_1\ dz_2|^2\frac{dy_1dt_1}{t_1^{n+1}}\frac{dy_2dt_2}{t_2^{m+1}}\\ \lesssim&\|a\|_2^2\int_{2l_I}^{\infty}\int_{\mathbb{R}^n}\int_I\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}\frac{|z_1-z_1'|^{2\alpha}}{(t_1+|y_1-z_1|)^{2n+2\alpha}}dz_1\frac{dy_1dt_1}{t_1^{n+1}}\\ &\times\int_0^{2l_J}\int_{\mathbb{R}^m}\int_J\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\frac{t_2^{2\beta}}{(t_2+|y_2-z_2|)^{2m+2\beta}}dz_2\frac{dy_2dt_2}{t_2^{m+1}}.\end{aligned}$$

Note that $2m+2\beta\leq 2m+2\varepsilon_2\leq m\lambda_2$. By Lemma 3.1, we obtain

$$\begin{aligned}(3.13)\qquad &\int_0^{2l_J}\int_{\mathbb{R}^m}\int_J\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\frac{t_2^{2\beta}}{(t_2+|y_2-z_2|)^{2m+2\beta}}dz_2\frac{dy_2dt_2}{t_2^{m+1}}\\ &\lesssim\int_0^{2l_J}\int_I\int_{\mathbb{R}^m}\frac{t_2^{2m+2\varepsilon_2+2\beta}}{\big(t_2+|x_2-y_2|\big)^{2m+2\varepsilon_2}}\frac{1}{(t_2+|y_2-z_2|)^{2m+2\beta}}dy_2\frac{dz_2dt_2}{t_2^{m+1}}\\ &\lesssim\int_0^{2l_J}\int_J\frac{t_2^{2\beta}}{\big(t_2+|x_2-z_2|\big)^{2m+2\beta}}\frac{dz_2dt_2}{t_2^{m+1}}\lesssim l_J^{m+2\beta}|x_2-z_2|^{-2m-2\beta}.\end{aligned}$$

Putting inequality (3.10) and (3.13) into inequality (3.12), we have

$$A_2 \lesssim l_I^{2\alpha} l_J^{2\beta} |x_1 - z_1|^{-2n-2\alpha} |x_2 - z_2|^{-2m-2\beta}.$$

Estimate for A_3. A_3 is symmetric with A_2, thus,

$$A_3 \lesssim l_I^{2\alpha} l_J^{2\beta} |x_1 - z_1|^{-2n-2\alpha} |x_2 - z_2|^{-2m-2\beta}.$$

Estimate for A_4. Similar to A_1 and A_2, the vanishing, support and size condition of rectangle atom, size condition of K_{t_1,t_2}, Hölder inequality and inequality (3.13) lead to

$$A_4 \lesssim l_I^{2\alpha} l_J^{2\beta} |x_1 - z_1|^{-2n-2\alpha} |x_2 - z_2|^{-2m-2\beta}.$$

Hence, whenever $x \in E_1$, we get

$$|g_{\vec{\lambda}}^*(a)(x)|^2 \le \sum_{i=1}^{4} A_i \lesssim l_I^{2\alpha} l_J^{2\beta} |x_1 - z_1|^{-2n-2\alpha} |x_2 - z_2|^{-2m-2\beta}.$$

This yields that

$$\iint_{E_1} |g_{\vec{\lambda}}^*(a)(x_1, x_2)| dx_1 dx_2 \lesssim \gamma^{-\alpha-\beta}.$$

Thus, we have proved inequality (2.5).

Now, we are in the position to prove inequality (2.6).

• **Proof of (2.6).** Recall that $E_2 = \{(x_1, x_2) \in \mathbb{R}^n \times \mathbb{R}^m : x_1 \in \gamma_1 I, x_2 \notin \gamma J\}$, we have

$$\begin{aligned} E_2 \subset & \{(x_1, x_2) \in \mathbb{R}^n \times \mathbb{R}^m : x_1 \in 2I, x_2 \notin \gamma J\} \\ & \cup \{(x_1, x_2) \in \mathbb{R}^n \times \mathbb{R}^m : x_1 \in \gamma_1 I \backslash 2I, x_2 \notin \gamma J\} =: E_{2,1} \cup E_{2,2}. \end{aligned}$$

Therefore,

$$\begin{aligned} \iint_{E_2} |g_{\vec{\lambda}}^*(a)(x_1, x_2)| dx_1 dx_2 \lesssim & \iint_{E_{2,1}} |g_{\vec{\lambda}}^*(a)(x_1, x_2)| dx_1 dx_2 \\ & + \iint_{E_{2,2}} |g_{\vec{\lambda}}^*(a)(x_1, x_2)| dx_1 dx_2. \end{aligned}$$

Since $E_{2,2} \subset \{(x_1, x_2) \in \mathbb{R}^n \times \mathbb{R}^m : x_1 \notin 2I, x_2 \notin \gamma J\}$, taking $\gamma_1 = 2$ and repeating the proof of (2.5), we may get

$$\iint_{E_{2,2}} |g_{\vec{\lambda}}^*(a)(x_1, x_2)| dx_1 dx_2 \lesssim \gamma^{-\alpha-\beta}.$$

So, to prove Theorem 1.3, we only need to show that there exists a $\delta > 0$ such that

$$\iint_{E_{2,1}} |g_{\vec{\lambda}}^*(a)(x_1, x_2)| dx_1 dx_2 \lesssim \gamma^{-\delta}. \tag{3.14}$$

We also need to split the domain of variable t_1 and t_2 as follows,

$$\begin{aligned} (t_1, t_2) \in \mathbb{R}^+ \times \mathbb{R}^+ \subset & \big([2l_I, \infty) \times [2l_J, \infty)\big) \cup \big([2l_I, \infty) \times (0, 2l_J)\big) \\ & \cup \big((0, 2l_I) \times [2l_J, \infty)\big) \cup \big((0, 2l_I) \times (0, 2l_J)\big) \\ & = F_1 \cup F_2 \cup F_3 \cup F_4. \end{aligned}$$

Hence, we may write

$$\begin{aligned}
&\iint_{E_{2,1}} |g^*_{\vec{\lambda}}(a)(x_1,x_2)|dx_1dx_2\\
&\lesssim \sum_{i=1}^{4}\iint_{E_{2,1}}\Big(\iint_{\mathbb{R}^{m+1}_+}\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\iint_{\mathbb{R}^{n+1}_+}\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}\mathbf{1}_{F_i}(t_1,t_2)\\
&\qquad\times\Big|\iint_{\mathbb{R}^n\times\mathbb{R}^m}K_{t_1,t_2}(y_1,y_2,z_1,z_2)a(z_1,z_2)dz_1\,dz_2\Big|^2\frac{dy_1dt_1}{t_1^{n+1}}\frac{dy_2dt_2}{t_2^{m+1}}\Big)^{1/2}dx_1dx_2\\
&=:\sum_{i=1}^{4}B_i.
\end{aligned}$$

Now, we need to give the estimate for each B_i.

Estimate for B_1. Since $x\in E_{2,1}$ and $(t_1,t_2)\in F_1$, then for any $z_1\in I$, $z_2\in J$, we have $t_1\geq 2l_I\geq 2|z_1-z_1'|$ and $t_2\geq 2l_J\geq 2|z_2-z_2'|$. Similar to inequality (3.9), by the vanishing, support and size condition of $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ rectangle atom, Hölder estimate of K_{t_1,t_2} and Hölder inequality, we have

(3.15)

$$\begin{aligned}
B_1\lesssim\|a\|_2^2\iint_{E_{2,1}}\Big(&\int_{2l_I}^{\infty}\int_{\mathbb{R}^n}\int_I\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}\frac{|z_1-z_1'|^{2\alpha}}{(t_1+|y_1-z_1|)^{2n+2\alpha}}dz_1\frac{dy_1dt_1}{t_1^{n+1}}\\
&\times\int_{2l_J}^{\infty}\int_{\mathbb{R}^m}\int_J\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\frac{|z_2-z_2'|^{2\beta}}{(t_2+|y_2-z_2|)^{2m+2\beta}}dz_2\frac{dy_2dt_2}{t_2^{m+1}}\Big)^{1/2}dx_1dx_2.
\end{aligned}$$

Since $x\in E_{2,1}$, then we have $t_1\geq 2|x_1-z_1|$. By Lemma 3.1, we obtain

$$\begin{aligned}
&\int_{2l_I}^{\infty}\int_{\mathbb{R}^n}\int_I\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}\frac{|z_1-z_1'|^{2\alpha}}{(t_1+|y_1-z_1|)^{2n+2\alpha}}dz_1\frac{dy_1dt_1}{t_1^{n+1}}\\
&\lesssim l_I^{2\alpha}\int_{2l_I}^{\infty}\int_I\int_{\mathbb{R}^n}\frac{t_1^{2n+2\varepsilon_1}}{\big(t_1+|x_1-y_1|\big)^{2n+2\varepsilon_1}}\frac{1}{(t_1+|y_1-z_1|)^{2n+2\alpha}}dy_1\frac{dz_1dt_1}{t_1^{n+1}}\\
&\lesssim l_I^{2\alpha}\int_{2l_I}^{\infty}\int_I\frac{t_1^n}{\big(t_1+|x_1-z_1|\big)^{2n+2\alpha}}\frac{dz_1dt_1}{t_1^{n+1}}\lesssim l_I^n.
\end{aligned}\tag{3.16}$$

Similarly, we may get

$$\begin{aligned}
&\int_{2l_J}^{\infty}\int_{\mathbb{R}^m}\int_J\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\frac{|z_2-z_2'|^{2\beta}}{(t_2+|y_2-z_2|)^{2n+2\beta}}dz_2\frac{dy_2dt_2}{t_2^{m+1}}\\
&\qquad\lesssim l_J^{m+2\beta}|x_2-z_2|^{-2m-2\beta}.
\end{aligned}\tag{3.17}$$

Putting the inequality (3.16), (3.17) into inequality (3.15), recall that $\|a\|_2^2\lesssim|R|^{-1}=l_I^{-n}l_J^{-m}$, we may arrive at $B_1\lesssim\gamma^{-\beta}$.

Estimate for B_2. Since $x\in E_{2,1}$ and $(t_1,t_2)\in F_2$, for any $z_1\in I$, we have $t_1\geq 2l_I\geq 2|z_1-z_1'|$. By the vanishing, support and size condition of rectangle atom, mixed Hölder

and size condition of K_{t_1,t_2} and Hölder inequality, similar as (3.12), we have

(3.18)

$$B_2 \lesssim \|a\|_2^2 \iint_{E_{2,1}} \Big(\int_{2l_I}^{\infty} \int_{\mathbb{R}^n} \int_I \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \frac{|z_1-z_1'|^{2\alpha}}{(t_1+|y_1-z_1|)^{2n+2\alpha}} dz_1 \frac{dy_1dt_1}{t_1^{n+1}} \\ \times \int_0^{2l_J} \int_{\mathbb{R}^m} \int_J \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \frac{t_2^{2\beta}}{(t_2+|y_2-z_2|)^{2m+2\beta}} dz_2 \frac{dy_2dt_2}{t_2^{m+1}} \Big)^{1/2} dx_1dx_2.$$

Putting inequalities (3.16) and (3.13) into (3.18), we may get $B_2 \lesssim \gamma^{-\beta}$.

Estimate for B_3. If $x \in E_{2,1}$, $t \in F_3$, then for any $z_2 \in J$, we have $t_2 \geq 2l_J \geq 2|z_2 - z_2'|$. The vanishing, support and size condition of $H^1(\mathbb{R}^n \times \mathbb{R}^m)$ rectangle atom, combination of Carleson and Hölder conditions of K_{t_1,t_2}, Hölder inequality and (3.13) imply that

$$B_3 = \iint_{E_{2,1}} \Big(\iint_{\mathbb{R}_+^{m+1}} \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \iint_{\mathbb{R}_+^{n+1}} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \mathbf{1}_{F_3}(t_1,t_2) \\ \times |\iint_{\mathbb{R}^n\times\mathbb{R}^m} (K_{t_1,t_2}(y,z) - K_{t_1,t_2}(y,(z_1,z_2')))a(z_1,z_2)dz_1dz_2|^2 \frac{dy_1dt_1}{t_1^{n+1}} \frac{dy_2dt_2}{t_2^{m+1}} \Big)^{\frac{1}{2}} dx_1dx_2$$

$$\lesssim \|a\|_{L^2} \iint_{2I\times(\gamma J)^c} \Big(\iint_{\mathbb{R}_+^{m+1}} \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \iint_{\mathbb{R}_+^{n+1}} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \mathbf{1}_{F_2}(t_1,t_2) \\ \iint_{\mathbb{R}^n\times\mathbb{R}^m} |K_{t_1,t_2}(y,z) - K_{t_1,t_2}(y,(z_1,z_2'))|^2 dz_1 \ dz_2 \frac{dy_1dt_1}{t_1^{n+1}} \frac{dy_2dt_2}{t_2^{m+1}} \Big)^{1/2} dx_1dx_2$$

$$\lesssim \|a\|_{L^2} |2I|^{1/2} \int_{(\gamma J)^c} \Big(\int_{2l_j}^{\infty} \int_{\mathbb{R}^m} \int_J \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} (\iint_{\widehat{I}} \int_{\mathbb{R}^n} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \\ |K_{t_1,t_2}(y,z) - K_{t_1,t_2}(y,(z_1,z_2'))|^2 dz_1 \frac{dy_1dx_1dt_1}{t_1^{n+1}}) \frac{dz_2dy_2dt_2}{t_2^{m+1}} \Big)^{1/2} dx_2$$

$$\lesssim \|a\|_{L^2} |I|^{\frac{1}{2}} \int_{(\gamma J)^c} \Big(\int_{2l_j}^{\infty} \int_{\mathbb{R}^m} \int_J \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \frac{|z_2-z_2'|^{\beta}}{(t_2+|y_2-z_2|)^{m+\beta}} \frac{dz_2dy_2dt_2}{t_2^{m+1}} \Big)^{\frac{1}{2}} dx_2$$

$$\lesssim \|a\|_{L^2} |I|^{\frac{1}{2}} l_J^{m/2+\beta} \int_{(\gamma J)^c} |x_2-z_2|^{-m-\beta} dx_2 \lesssim \gamma^{-\beta}.$$

Estimate for B_4. Similarly, we have

$$B_4 = \iint_{E_{2,1}} \Big(\iint_{\mathbb{R}_+^{m+1}} \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \iint_{\mathbb{R}_+^{n+1}} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \mathbf{1}_{F_4}(t_1,t_2) \\ |\iint_{\mathbb{R}^n\times\mathbb{R}^m} K_{t_1,t_2}(y,z)a(z_1,z_2)dz_1 \ dz_2|^2 \frac{dy_1dt_1}{t_1^{n+1}} \frac{dy_2dt_2}{t_2^{m+1}} \Big)^{1/2} dx_1dx_2$$

$$\lesssim \|a\|_{L^2} \iint_{2I\times(\gamma J)^c} \Big(\iint_{\mathbb{R}_+^{m+1}} \Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2} \iint_{\mathbb{R}_+^{n+1}} \Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1} \mathbf{1}_{F_4}(t_1,t_2) \\ \iint_{\mathbb{R}^n\times\mathbb{R}^m} |K_{t_1,t_2}(y,z)|^2 dz_1 \ dz_2 \frac{dy_1dt_1}{t_1^{n+1}} \frac{dy_2dt_2}{t_2^{m+1}} \Big)^{1/2} dx_1dx_2$$

$$\lesssim \|a\|_{L^2}(2I)^{1/2}\int_{(\gamma J)^c}\Big(\int_0^{2l_j}\int_{\mathbb{R}^m}\int_J\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\Big(\iint_{\widehat{I}}\int_{\mathbb{R}^n}\Big(\frac{t_1}{t_1+|x_1-y_1|}\Big)^{n\lambda_1}$$
$$|K_{t_1,t_2}(y,z)|^2dz_1\frac{dy_1dx_1dt_1}{t_1^{n+1}}\Big)\frac{dz_2dy_2dt_2}{t_2^{m+1}}\Big)^{1/2}dx_2$$
$$\lesssim \|a\|_{L^2}|I|^{\frac{1}{2}}\int_{(\gamma J)^c}\Big(\int_{2l_j}^{\infty}\int_{\mathbb{R}^m}\int_J\Big(\frac{t_2}{t_2+|x_2-y_2|}\Big)^{m\lambda_2}\frac{t_2^{\beta}}{(t_2+|y_2-z_2|)^{m+\beta}}\frac{dz_2dy_2dt_2}{t_2^{m+1}}\Big)^{\frac{1}{2}}dx_2$$
$$\lesssim \|a\|_{L^2}(2I)^{1/2}l_J^{m/2+\beta}\int_{(\gamma J)^c}|x_2-z_2|^{-m-\beta}dx_2\lesssim\gamma^{-\beta}.$$

Combining the estimate for $B_i(i=1\cdots 4)$, we get

$$\iint_{E_{2,1}}|g_\lambda^*(a)(x_1,x_2)|dx_1dx_2\lesssim\gamma^{-\beta}.$$

This completes the proof of (3.14). Since we have reduced the proof of Theorem 1.2 to inequality (3.14), the proof of Theorem 1.2 is finished. □

4. Proof of the corollaries

We need the following lemma:

Theorem C. (*Calderón-Zygmund Lemma* [3]) *Let $\alpha>0$ be given and $f\in L^p(\mathbb{R}^n\times\mathbb{R}^m)$, $1<p<2$. Then we may write $f=f_1+f_2$ where $f_1\in L^2(\mathbb{R}^n\times\mathbb{R}^m)$ and $f_2\in H^1(\mathbb{R}^n\times\mathbb{R}^m)$ with $\|f_1\|_2^2\le\alpha^{2-p}\|f\|_{L^p_p}$; and $\|f_2\|_{H^1}\le c\alpha^{1-p}\|f\|_{L^p_p}$, where c is a universal constant.*

A trivial corollary of Theorem C will lead to the following result.

Lemma 4.1. *Let T be a sublinear operator which is bounded from $H^1(\mathbb{R}^n\times\mathbb{R}^m)$ to $L^1(\mathbb{R}^n\times\mathbb{R}^m)$ and bounded on $L^2(\mathbb{R}^n\times\mathbb{R}^m)$. Then T is bounded on $L^p(\mathbb{R}^n\times\mathbb{R}^m)$ for all $1<p<2$.*

Thus, the L^p boundedness in Corollary 1 and Corollary 2 follows from the interpolation between (H^1,L^1) and (L^2,L^2).

References

[1] M. Cao, Q. Xue, *On the boundedness of bi-parameter Littlewood-Paley g_λ^*-function*, availiable at http://arxiv.org/abs/1512.00569.

[2] S. Y. Chang, R. Fefferman, *A continuous version of duality of H^1 with BMO on bi-disc*, Ann. of Math., **112** (1980), 179-201.

[3] S. Y. Chang, R. Fefferman, *The Calderón-Zygmund decomposition on product domains*, Amer. J. Math., **104** (1982), 455-468.

[4] R. Fefferman, *Calderón-Zygmund theory for product domain-H^p theory*, Proc. Nat. Acad. Sci. USA, **83** (1986), 840-843.

[5] R. Fefferman, *Harmonic analysis on product spaces*, Ann. of Math., **126** (1987), 109-130.

[6] R. Fefferman, E. Stein, *Singular integrals on product spaces*, Adv. Math., **45** (2) (1982), 117-143.

[7] T. Hytönen, *Representation of singular integrals by dyadic operators, and the A_2 theorem*, preprint (2011).

[8] T. Hytönen, H. Martikainen, *Non-homogeneous $T1$ theorem for bi-parameter singular integrals*, Adv. Math., **261** (2014), 220-273.

[9] J.-L. Journé, *Calderón-Zygmund operators on product spaces*, Rev. Mat. Iberoamericana., **1** no. 3 (1985), 55-91.
[10] H. Martikainen, *Representation of bi-parameter singular integrals by dyadic operators*, Adv. Math., **229** (3) (2012), 1734-1761.
[11] Y. Ou, *A $T(b)$ Theorem on product spaces*, Trans. Amer. Math. Soc., **367** (2015), 6159-6197.
[12] S. Pott and P. Villarroya, *A $T(1)$ theorem on product spaces*, preprint (2013).
[13] R.F. Gundy, E.M. Stein, *H^p theory for the poly-disc*, Proc. Nat. Acad. Sci. USA, **76** (1979), 1026-1029.
[14] Y. Han, M.Y. Lin, C. Lin, Y. Lin, *Calderón-Zygmund operators on product Hardy spaces*, J. Funct. Anal., **258** (2010), 2834-2861.
[15] Y. Han, D. Yang, *H^p boundedness of Calderón-Zygmund operators on product spaces*, Math. Z., **249** (2005), 869-881.
[16] J.L Journé, *Calderón-Zygmund operators on product spaces*, Rev. Mat. Iberoamericana, **1** (1985), 55-91.
[17] M.P. Malliavin, P. Malliavin, *Intégrales de Lusin-Calderón pour les fonctions biharmoniques*, Bull. Sci. Mathe., **101** (1977), 357-384.
[18] H. Martikainen, *Boundedness of a class of bi-parameter square function in the upper half-space*, J. Funct. Anal., **267** (2014), 3580-3597.
[19] Q. Xue, J. Yan, *On multilinear square function and its applications to multilinear Littlewood-Paley operators with non-convolution type kernels*, J. Math. Anal. Appl., **422** (2015), 535-559.
[20] D. Yang, Y. Zhou, *Boundedness of Marcinkiewicz integrals and their commutators in $H^1(\mathbb{R}^n \times \mathbb{R}^m)$*, Sci. China, **Series A 49** (2006), 770-790.

Zhengyang Li, School of Mathematical Sciences, Beijing Normal University, Laboratory of Mathematics and Complex Systems, Ministry of Education, Beijing 100875, People's Republic of China

E-mail address: zhengyli@mail.bnu.edu.cn

Qingying Xue, School of Mathematical Sciences, Beijing Normal University, Laboratory of Mathematics and Complex Systems, Ministry of Education, Beijing 100875, People's Republic of China

E-mail address: qyxue@bnu.edu.cn